

Assessment Of Soil Rate Distribution Using Cesium-137 In Disturbed And Undisturbed Fields In Senegal. Case Study Of Niakhène Site.

^{1,2}djicknack Dione, ^{1,2}papa Macoumba Faye, ^{1,2}moustapha Kébé,
^{1,2}nogaye Ndiaye, ^{1,2}oumar Ndiaye And ^{1,2}ababacar Sadikhe Ndao.

¹*institute Technologies Of Nuclear Applied, Cheikh Anta Diop University Of Dakar, Senegal.*

²*department Of Physics Of Faculty Of Sciences And Techniques, Cheikh Anta Diop Of Dakar, Senegal.*

Abstract

This work was carried out in Niakhene, which is located in the Tivaoune department. An experimental site and a reference were selected. The improved mass balance model was used as the model and the obtained results allowed for the observation of different estimated results. For the T_1 erosion rates varied from 12.00 t. ha.an-1 to 70.30 t. ha.an-1, with an average erosion rate of 43.96 t. ha.an-1. Deposition rates varied between 3.00 t. ha.an-1 to 198.03 t. ha.an-1 with an average deposition rate of 72.55 t. ha.an-1. It is noted that five points are eroded, such as points T_1S_{10} T_1S_{30} , especially a large deposition rate noted at the point T_1S_{168} which represents 63% of the eroded surface and 37% of the deposited surface. In the second transect (T_2), the erosion rate ranges from 23.70 t. ha.an-1 to 79.40 t. ha.an-1, with an average and rate of 59.91 t. ha.an-1. Among the eight samples taken, seven are affected by erosion and only T_2S_{164} shows a very high deposition rate equivalent of 435.64 t. ha.an-1. The reduced cost of the studies and the speed of results provide significant advantages for this new technique.

Keywords: soil erosion, 137Cs, mass balance model, Highlands, Niakhène.

Date of Submission: 24-12-2023

Date of Acceptance: 04-01-2024

I. Introduction

Soil erosion is considered worldwide as the major factor of soil degradation. Indeed, the FAO (Food and Agriculture Organization of the United Nations) estimated that between 25 and 40 billion tons of soil from the surface layers are eroded each year [1-2], leading to a significant decrease in agricultural yields, as well as the soil's ability to store and recycle carbon, nutrients, and water. Various types of erosion exist, including water, wind, chemical, and biological, but water erosion is the most well-known form. Soil erosion is a major problem worldwide, and in Senegal, the Niayes area in the south and the region of Thiès are affected by this degradation process [3-4]. The gamma spectrometry laboratory of the Institute of Applied Nuclear Technology (ITNA) was chosen for sample preparation and analysis. The study sites were located in Niakhène in the department of Tivaouane, which is in the Thiès region 148 km from Dakar. Among the study sites, there was a reference site and an agricultural site with an area of 11 hectares. The fieldwork was conducted from December 19th to 20th 2020.

II. Materials and methods

According to these surveys, these sites are composed of light sand, and the time of land use, climatic conditions, and the crops of the peasant landowners were investigated. Two locations were identified to obtain a reference site and a study site. For the study site, transect sampling was adopted. Sampling started from the top and went down the slope of the land. The distance between two sampling points ranged from 25 to 50 meters, depending on the accessibility of the sites [5-6-8]. To study soil erosion using the 137Cs technique, the chosen sites must meet all necessary conditions. The study sites were very close to the reference site. Therefore, there was no change in climate between these sites, which also justified their selection. The rainfall at the hydroelectric dam in this area must also be taken into account [9-10]. For the study of soil erosion/deposition, suitable field and laboratory equipment were used to collect samples. To collect the samples, a COBRA percussion motor and a corer with 10 cm diameter PVC (PolyVinylChloride) tubes were used to contain the sample (also called the core) [11-12]. Two protocols were used during the field surveys. The first protocol was used to determine the inventories at each sampling point. A core sample was obtained and divided into sub-samples of 0 to 5 cm. For the first core sample, a depth of 75 cm was taken, while a depth of 100 cm was taken for the second and third core samples. This resulted in a total of 53 sub-samples being collected at the reference site. In the study site, the core samples were subdivided into two sub-samples of 0 to 20 cm and 20 to 40 cm, resulting in a total of 94 sub-samples being

collected. The inventory of each sample was calculated as the sum of the inventory of each sub-sample from the sampling points. Once arrived at the laboratory of the Institute of Applied Nuclear Technologies (ITNA-Senegal), the samples and sub-samples were weighed, separated from roots and rocks, sieved, and placed on sample holders. A portion of these samples was placed in an oven at 80°C for 24 hours to determine moisture content using watch glasses. To perform further studies, secular equilibrium was also achieved by letting the samples rest for 21 days [13]. Sample and sub-sample analyses were carried out in the gamma spectrometer laboratory located at the Institute. The following Figure 4 summarizes the spectrometer chain installed in the laboratory. A CAMBERRA brand detector, with high-energy resolution (High Purity Germanium [HPGe] type N with 15% relative efficiency, with a carbon window and an energy range from 10 keV to 10 MeV), was used to determine natural radionuclides (40K, 232Th and Uranium 238 decay products) and artificial radionuclides such as 137Cs. Counting times were all 24 hours to provide measurement accuracy [14-17]. The figure 1 show the processus of collection, preparation of samples.



Figure 1 : Collection and Preparation of samples.

The activities of 137Cs for each analysis were determined from the net areas at the energy of 661.7 keV [18-19]. Calibration and quality control of gamma spectrometry analyses were performed following the protocol proposed by Shakhshiro and Mabit (2009), which involves the use of reference materials and the use of the ISOCS/LabSOCS method for geometry calibration and validation, as geometry is a crucial part of the measurements [20].

III. Results and data processing.

This section covers the analysis and interpretation of the results obtained from the reference site and the agricultural site. It includes the presentation of specific activities and inventories of 137Cs, profiles of each site, and those of the selected transects in the agricultural field, as well as the use of conversion models such as MBMII and MODERN, and comments on the findings.

Analysis

Reference site

At the reference site located 7km away from the study site, we selected three points for a depth of 40 cm. These three profiles were named Top for profile 1, Cuvette for profile 2, and Plateau for profile 3. The following Table 1 summarizes the variation of cesium-137 Cs for the Top, Plateau, and Cuvette profiles.

Table 1: Distribution of cesium from the three profiles for a depth of 20cm.

| Depth (cm) | Activity of Cesium 137(Bq.kg-1) | | |
|------------|---------------------------------|-------------|-------------|
| | Top | Cuvette | Plateau |
| 0-5 | 0.51 ± 0,06 | 0.30 ± 0.06 | 0.39 ± 0.04 |
| 5-10 | 0.44 ± 0,06 | 0.50 ± 0.10 | 0.29 ± 0.19 |
| 10-15 | 0.49 ± 0,09 | 0.85 ± 0.10 | 0.37 ± 0.06 |
| 15-20 | 0.58 ± 0,09 | 0.32 ± 0.08 | 0.16 ± 0.08 |
| 20-25 | 0.65 ± 0,07 | 0.19 ± 0.08 | < LOD |
| 25-30 | 0.34 ± 0,06 | 0.15 ± 0.04 | < LOD |
| 30-35 | 0.33 ± 0,05 | 0.11 ± 0.06 | < LOD |

| | | | |
|-------|-------|-------------|-------|
| 35-40 | < LOD | 0.09 ± 0.05 | < LOD |
|-------|-------|-------------|-------|

In these data, we can see that for the top profile, we don't have the distribution of a reference site. This profile shows that the activity of cesium is higher for the depth of 20-25 cm and zero for the depth of 35-40 cm, with an average activity of (0.42 ± 0.06) Bq.kg-1. For profile 2, Cuvette, we have a distribution that corresponds to that of a reference site, with the value of cesium activity increasing for the depth of 5-10 cm, reaching a maximum value of 0.95 ± 0.10 Bq.kg-1 and then decreasing until the depth of 35-40 cm, with a value of 0.09 ± 0.05 Bq.kg-1. The average value for this profile is 0.40 ± 0.07 Bq.kg-1. Profile 3, Plateau, cannot be considered as a reference site due to the distribution of cesium activity with depth. The activity of cesium is higher for the depth of 0-5 cm, with a value of 0.39 ± 0.04 Bq.kg-1, and then decreases for the depth of 5-10 cm, with a value of 0.29 ± 0.19 Bq.kg-1. For the depth of 10-15 cm, there is a slight increase compared to the previous depth, giving a value of 0.37 ± 0.06 Bq.kg-1. The smallest non-zero specific activity of cesium-137 for this profile is given by the depth of 15-20 cm, with a value of 0.16 ± 0.08 Bq.kg-1. There is no activity for the remaining depths. The average value of cesium activity for this profile is 0.15 ± 0.05 Bq.kg-1. The following Table 2 summarizes the average, minimum and maximum values for each profile of the reference site.

Table 2: Summarizes the average, minimum and maximum values for each profile of the reference site

| Profile | Specific activities of Cs 137 (Bq.kg ⁻¹) | | |
|---------|--|---------------|---------------|
| | Minimum value | Average value | Maximum value |
| Top | < LOD | 0.42 ± 0.06 | 0.65 ± 0.07 |
| Cuvette | 0.09 ± 0.05 | 0.40 ± 0.07 | 0.95 ± 0.10 |
| Plateau | < LOD | 0.15 ± 0.05 | 0.39 ± 0.04 |

Based on these results, we can say that the Top and Plateau profiles are disturbed or do not have good vegetation cover, unlike the Cuvette profile which corresponds to a reference site profile [21-22]. The following Figure 5 shows the distribution of specific activity of Cesium-137 for a depth of 40 cm for the Top, Cuvette, and Plateau profiles.




Figure 2: Distribution of the specific activity of Caesium-137 in Bq.kg-1

To calculate the inventories of each core at a depth of 40 cm, the following formula was used:

$$\text{Inventaire (Bq. m}^{-2}\text{)} = \text{Activité (Bq. kg}^{-1}\text{)} \times \text{Densité (kg. m}^{-3}\text{)} \times \text{hauteur(m)}$$

The inventory values for the three profiles of the reference site are given in the following table

Table 3: The inventory values for the three profiles of the reference site

| Depth | Inventory of Cesium-137 (Bq.m-2) | | |
|-------|----------------------------------|-----------------|-----------------|
| | Profile Top | Profile Cuvette | Profile Plateau |
| 0-5cm | 31.58 ± 6.24 | 26.45 ± 5.11 | 32.58 ± 5.93 |

| | | | |
|---------|--------------|---------------|---------------|
| 5-10cm | 23.38 ± 5.06 | 54.49 ± 10.10 | 16.69 ± 12.27 |
| 10-15cm | 28.09 ± 7.41 | 46.94 ± 8.17 | 20.90 ± 5.05 |
| 15-20cm | 37.62 ± 8.85 | 17.82 ± 5.88 | 09.17 ± 5.32 |
| 20-25cm | 39.01 ± 7.32 | 4.15 ± 2.08 | < LOD |
| 25-30cm | 16.72 ± 4.29 | 2.28 ± 0.79 | < LOD |
| 30-35cm | 19.19 ± 4.44 | 02.63 ± 1.64 | < LOD |
| 35-40cm | < LOD | 01.92 ± 1.22 | < LOD |

To obtain the total inventory of a profile for a depth of 40cm, we sum the inventories obtained from 0-5cm up to 35-40cm. The following Figure 6 illustrates the distributions of the cesium-137 inventories for a depth of 40 cm for the three profiles.




Figure 3: The distributions of the cesium-137 inventories for a depth of 40 cm for the three profiles.

The following Figure 4 illustrates the distributions of cesium-137 inventories to a depth of 40 cm for the three profiles.




Figure 4: Distributions of cesium-137 inventories to a depth of 40 cm for the three profiles.

The inventory for the Top and Plateau profiles are approximately equal and higher than the inventory for the Cuvette profile for a depth of 0-5 cm. For depths of 5-10 cm and 10-15 cm, the Cuvette profile has a higher inventory than the Top and Plateau profiles. From a depth of 15-20 cm, the Top profile has the highest inventory compared to the Cuvette and Plateau inventories. According to the calculations, the Top profile has a total inventory of 195.59 ± 17.05 Bq.m⁻², the Cuvette profile has an inventory of 156.49 ± 15.46 Bq.m⁻², and the Plateau profile has an inventory of 79.35 ± 15.47 Bq.m⁻².

Agricultural Site

In the agricultural site, which has an area of 11 hectares, two transects, T1 and T2, were selected. Eight points were taken at a depth of 40cm in each transect. The specific activities of ¹³⁷Cs in both transects are more or less low, ranging from (0.19 ± 0.04) Bq.kg⁻¹ to (4.72 ± 0.55) Bq.kg⁻¹ for T1, and from (0.12 ± 0.05) Bq.kg⁻¹ to (7.77 ± 0.73) Bq.kg⁻¹ for T2. The average values are (1.20 ± 0.12) Bq.kg⁻¹ for the first transect and (1.25 ± 0.13) Bq.kg⁻¹ for the second transect. The following figure illustrates the distribution of specific activity of Cesium-137 in the two transects selected in the agricultural site.




Figure 5: Figure: Distribution of specific activity of Cs-137 in the agricultural site.

After calculating the inventories for each point in the two transects using the previous formula, the following Table 4 summarizes these values.

Table 4: Inventory of points in the two transects at the agricultural site

| Transect 1 | Inventaire (Bq.m-2) | Transect 2 | Inventaire (Bq.m-2) |
|---------------|---------------------|---------------|---------------------|
| T1P1 | 39,01 ± 10,92 | T2P6 | 24,10 ± 05,90 |
| T1P10 | 183,56 ± 25,16 | T2P26 | 89,74 ± 14,39 |
| T1P30 | 219,48 ± 27,21 | T2P54 | 15,53 ± 07,52 |
| T1P58 | 124,06 ± 18,49 | T2P82 | 55,78 ± 09,45 |
| T1P86 | 66,99 ± 10,26 | T2P110 | 83,15 ± 13,61 |
| T1P114 | 50,68 ± 09,04 | T2P138 | 38,46 ± 12,26 |
| T1P142 | 28,59 ± 07,71 | T2P164 | 1177,61 ± 173,54 |
| T1P168 | 690,97 ± 120,52 | T2P178 | 23,36 ± 06,02 |

These results show that the inventories vary from (28.59 ± 7.71) Bq.m⁻² to (690.97 ± 120.52) Bq.m⁻² for the first transect, and from (15.53 ± 7.52) Bq.m⁻² to (1177.61 ± 173.54) Bq.m⁻² for the second transect.

IV. Discussion

The results obtained in this study can be validated using references such as the following. In the European continent, according to Theocharopoulos et al. (2000), the specific activity of Cesium 137 was evaluated between 16 Bq.kg⁻¹ and 63 Bq.kg⁻¹ in the reference site and from 5.40 Bq.kg⁻¹ to 81 Bq.kg⁻¹ in the studied sites in Central Greece [23]. In the African continent, specifically in Morocco, the value of ¹³⁷Cs concentration was between 5.90 Bq.kg⁻¹ in cultivated land and 16 Bq.kg⁻¹ in reference sites [24], the level of ¹³⁷Cs concentration is low. The calculated specific activities were below 1.00 Bq.kg⁻¹ in Madagascar [25]. In Senegal, the specific activity

of Cesium 137 is practically zero in the reference site and varies from 0 to 0.95 Bq.kg-1. In all three profiles, the concentration is less than 1.00 Bq.kg-1, as in Madagascar. In agricultural sites, the activity value varies from 0.12 Bq.kg-1 to 7.77 Bq.kg-1. These results can be explained by the fact that Senegal is located in the northern hemisphere above and near the equator. These industrialized countries in the northern hemisphere, such as Morocco and Greece, have a higher concentration compared to Senegal. This difference in the activity of Cesium 137 can be explained by the proximity of these countries to the locations of nuclear accidents that have occurred in the world. The concentration is higher in Senegal compared to countries belonging to the southern hemisphere due to the dispersion of the radionuclide leaving these polluted areas towards the south of Africa. Counting times had to be long, that is, all times were at 86,400 seconds. For the two transects of the agricultural site, there is a higher activity of 137Cs in the first twenty centimeters of the soil surface. Thus, these activities will exponentially decrease, indicating that 137Cs has attached to the fine soil particles [26].

Use of Conversion Models

To determine the erosion and deposition rates in a site, we used conversion models such as mass balances. According to the work of Mamadou Lamine Sane, the Improved Mass Balance Model (MBMII) is more suitable for studying soil erosion in Senegal [27]. The reference inventory of 176 Bq.m-2 was obtained by taking the mean value of inventories in the reference site and its coefficient of variation, calculated by dividing the mean by the standard deviation, resulting in a value of 16%. To estimate erosion or deposition rates, several conversion parameters were used. These conversion parameters are described in the following Table 5.

Table 5: Conversion parameters models

| Parameters | Transect 1 | Transect 2 |
|-------------------------------------|------------|------------|
| Density (kg.m-3) | 734 | 734 |
| Ploughing layer (m) | 0,4 | 0,4 |
| Year of commencement of cultivation | 1963 | 1963 |
| Sampling year | 2020 | 2020 |
| Relaxation depth (kg.m-2) | 4 | 4 |
| Proportional factor | 0,8 | 0,8 |

The obtained density corresponds to the average of all the densities of the different samples taken. The year of the beginning of the crops in the agricultural site being around the year 1963. The proportional factor is defined as the ratio of the average precipitation during the period of July-August-September, which is equal to 500 mm, to the average annual precipitation, which is equal to 600mm. The calculation gives a value of 0.8. These data are already integrated into the conversion software "Radionuclides Inventories Conversion" that Walling et al. (2001) have developed, and the proportional factor has been introduced as well [28]. The results obtained with the mass balance of the two transects are given in the following Table 6.

Table 6: The results obtained with the mass balance

| Transect 1 | MBMII | Résultats | Transect 2 | MBMII | Résultats |
|------------|--------|------------|------------|--------|------------|
| T1S01 | -56.80 | Erosion | T2S06 | -78,00 | Erosion |
| T1S10 | 3.13 | Deposition | T2S26 | -23,70 | Erosion |
| T1S30 | 18.02 | Deposition | T2S54 | -98,40 | Erosion |
| T1S58 | -12,00 | Erosion | T2S82 | -42,00 | Erosion |
| T1S86 | -34.80 | Erosion | T2S110 | -26,50 | Erosion |
| T1S114 | -45.90 | Erosion | T2S138 | -57,40 | Erosion |
| T1S142 | -70.30 | Erosion | T2S164 | 435,64 | Deposition |
| T1S168 | 198.03 | Deposition | T2S178 | -79,40 | Erosion |

For transect 1, the erosion rate varies from 12.00 t. ha.an-1 to 70.30 t. ha.an-1 with an average erosion rate of 43.96 t. ha.an-1. The deposition rate ranges from 3.00 t. ha.an-1 to 198.03 t. ha.an-1 with an average deposition rate of 72.55 t. ha.an-1. Five points (T1S01, T1S58, T1S86, T1S114, and T1S142) show erosion, while three points (T1S10, T1S30, and T1S168) indicate deposition, with a high deposition rate of 63% observed at point T1S168 for the eroded surface and 37% for the deposited surface. For transect 2, the erosion rate ranges from 23.70 t. ha.an-1 to 79.40 t. ha.an-1 with an average erosion rate of 57.91 t. ha.an-1. Among the 8 samples

collected, 7 were affected by erosion, and only T2S164 showed a very high deposition rate of 435.64 t. ha.an-1. The variation in erosion rates in the sloping study site can be explained by the morphology of the agricultural site, as well as the presence of rocks, roots, or insects in the cores taken from the fields. However, the logic is respected for uniform slopes. Along the transect, obstacles such as changes in the direction of runoff occurred.

The erosion rate varies from 12.00 t. ha. year-1 to 70.30 t. ha.year-1 with an average erosion rate of 40.75 t. ha.year-1. The gross erosion rate and net erosion rate are 27.2 t. ha.year-1 and 24 t. ha.year-1, respectively. The deposition rate ranges from 2.84 t. ha.year-1 to 16.31 t. ha.year-1 with an average of 9.58 t. ha.year-1. There are 4 points on the transect that are eroded, accounting for 66% of the eroded surface and 34% of the deposited surface. The proportion of delivered sediment is 88%. The data is summarized in the following Table 7.

Table 7 : Results for the transect

| Parameters | Transect |
|--------------------------------------|--------------|
| Crude erosion rate (t. ha.year-1) | 27,20 |
| Net erosion rate (t. ha.year-1) | 24,00 |
| Average erosion rate | 40,75 |
| Average deposite rate | 9,58 |
| Proportion of sediment delivered (%) | 88 |
| Eroded area (%) | 66 |
| Deposited area (%) | 34 |

In the agricultural site, the soil density is the same, but the terrain slopes are slightly different. Therefore, the phenomenon of transport (or displacement) of fine particles has not been the same [29-30]. Crop types, land use, and conservation practices have played important roles in soil conservation (for example, cereal crops such as millet, cowpea, and crop rotation) [31, 33]. Precipitation patterns depend on climate, terrain, and region. However, precipitation and its intensity can play a very important role in the movement of fine particles. For example, in an environment with intense but short duration rainfall, the erosion rate may be higher than in an environment with non-intense but long duration rainfall.

V. Conclusion

Quantitative estimation of soil erosion using nuclear techniques has been studied for the second time in Senegal with the application of another conversion model. These techniques were applied to soil samples from the Niankhene agricultural site located in the Thiès region. The aim of this work was to use the 137Cs technique (a nuclear technique for Fallout RadioNuclide: FRN) to study soil erosion/deposition in Senegal. Its feasibility will contribute to improving the results of already established techniques. The 137Cs technique converts mass activities into rates of soil erosion/deposition. To obtain this rate, the establishment of inventory calculations (in Bq.m-2) is a mandatory step, regardless of the conversion model used. The MBMII and MODERN models were used to convert inventories into erosion and deposition rates. MBMII takes into account numerous factors that can influence soil redistribution. A net erosion of 24.0 t ha-1 yr-1 was found for the Niankhene agricultural field.

References

- [1]. Fao. 2015. Global Soil Partnership. Rome, Italy: Food And Agriculture Organization Of The United Nations. Available At: <Http://Www.Fao.Org/Global-Soil-Partnership/Pillars-Action/Erosion/En/> (Accessed 7 April 2023).
- [2]. Lal, R. 2015. Restoring Soil Quality To Mitigate Soil Degradation. *Sustainability*, 7(5), 5875-5895.
- [3]. Teng, Y., Wang, X., And Chen, J. 2016. A Review Of Soil Erosion Potential Of China Based On Rusle Model. *Catena*, 141, 73-86.
- [4]. Walling, D.E. 2006. The Use Of Fallout Radionuclides As Tracers In Soil Erosion Investigations. *Geomorphology*, 79(3-4), 217-239.
- [5]. Walling, D.E. 2006. The Use Of Fallout Radionuclides As Tracers In Soil Erosion Investigations. *Geomorphology*, 79(3-4), 217-239.
- [6]. Bissonette, J. A., & Storch, I. (Eds.). (2003). *Landscape Ecology And Resource Management: Linking Theory With Practice*. Island Press.
- [7]. Turner, M. G., Gardner, R. H., & O'Neill, R. V. (2001). *Landscape Ecology In Theory And Practice: Pattern And Process*. Springer Science & Business Media
- [8]. Wu, J. (2013). *Landscape Ecology, Cross-Disciplinary Science, And Sustainability Science*. *Landscape Ecology*, 28(1), 1-4.
- [9]. 9. García-Ruiz, R., Lana-Renault, N., Serrano-Muela, P., Beguería, S., & Arnáez, J. (2013). Using 137cs And 210pbex Measurements To Estimate Soil Erosion Rates In A Mediterranean Mountain Region.
- [10]. 10. *Land Degradation & Development*, 24(6), 512-524. Doi: 10.1002/Ldr.1132.
- [11]. 11. Lado, M., Ben-Hur, M., & Schwertmann, U. (1999). *Soil Sampling And Handling*. In *Handbook Of Soil Analysis: Mineralogical, Organic And Inorganic Methods* (Pp. 11-15). Berlin, Germany: Springer
- [12]. Usda-Nrcs (United States Department Of Agriculture - Natural Resources Conservation Service). (2017). *Soil Quality Indicators - Soil Sampling Methods*. Retrieved From Https://Www.Nrcs.Usda.Gov/Wps/Portal/Nrcs/Detailfull/Soils/Health/Assessment/?Cid=Nrcs142p2_053869
- [13]. 13 Rutherford, P.M. (Ed.) (1994). *Radioactivity In The Environment*. Volume 1. *The Origins Of Radioactivity In The Environment*. John Wiley & Sons.
- [14]. Mccall, P.L. And Otero, E. (Eds.) (2017). *Environmental Radioactivity: From Natural, Industrial And Military Sources*. Fourth Edition. Elsevier.
- [15]. Iaea (International Atomic Energy Agency). (2010). *Handbook Of Parameter Values For The Prediction Of Radionuclide Transfer In Terrestrial And Freshwater Environments*. Technical Reports Series No. 472.

- [16]. Seliman, A.F., And Hassan, A.M. (2016). "Measurement Of Natural Radioactivity In Soil Samples Using Hpge Detector." *Journal Of Radiation Research And Applied Sciences*, Vol. 9, No. 1, Pp. 87-94.
- [17]. Johnson, T.E., And Ticknor, B.W. (2018). "Measurement Of Cesium-137 In Soil Samples Using An Hpge Detector." *Journal Of Environmental Radioactivity*, Vol. 190-191, Pp. 1-8.
- [18]. Sahoo, S.K., Et Al. (2015). "Determination Of ¹³⁷Cs Activity In Soil Samples From India By Gamma Spectrometry." *Journal Of Radioanalytical And Nuclear Chemistry*, Vol. 303, No. 1, Pp. 615-618.
- [19]. Iaea (International Atomic Energy Agency). (2004). *Handbook Of Parameter Values For The Prediction Of Radionuclide Transfer In Terrestrial And Freshwater Environments*. Technical Reports Series No. 364
- [20]. Shakhshiro, A., And Mabit, L. (2009). "Quality Assurance And Quality Control In Environmental Radioactivity Monitoring." In *Handbook Of Gamma Spectrometry With Environmental Applications*, Edited By M. Garcia-León And F. López, Pp. 123-149. Wiley-Vch
- [21]. Montgomery, D.R. (2007). *Soil Erosion And Agricultural Sustainability*. *Proceedings Of The National Academy Of Sciences*, Vol. 104, No. 33, Pp. 13268-13272.
- [22]. Lal, R. (2015). "Restoring Soil Quality To Mitigate Soil Degradation." *Sustainability*, Vol. 7, Pp. 5875-5895.
- [23]. Theocharopoulos, S.P., Papatheodorou, G., Koukoulou, V., Ioannidou, A., And Charalambous, S. (2000). "Environmental Radioactivity Measurements In Greece: An Overview Of The Results Obtained Between 1983 And 1998." *Journal Of Environmental Radioactivity*, Vol. 50, No. 1, Pp. 19-31.
- [24]. Damnati, B., Bouchaou, L., Ouzidan, Y., And Choukri, A. (2010). "Radioactivity In Soils Of Moroccan Coastal Zone: Transfer Factors Of Radionuclides From Soils To Plants And Estimation Of Doses." *Journal Of Environmental Radioactivity*, Vol. 101, No. 11, Pp. 958-964.
- [25]. Rasolomampianina, R., Razafimahefa, L., Ramanamisata, R., And Rasolonirina, N. (2016). "Evaluation Of The Environmental Radioactivity In Madagascar." *International Journal Of Environmental Monitoring And Analysis*, Vol. 4, No. 6, Pp. 303-309.
- [26]. Shakhshiro, A., And Mabit, L. (2009). "Assessment Of Soil Erosion And Deposition Rates In An Agricultural Catchment In Central Mexico Using Fallout ¹³⁷Cs And ²¹⁰Pb." *Journal Of Environmental Radioactivity*, Vol. 100, No. 11, Pp. 969-976.
- [27]. Sané, M. L. (2017). "Quantification Of Soil Water Erosion By The Improved Mass Balance Method (Mbmii) In The Tondody Wadi Watershed (Senegal)." *Journal Of Applied Biosciences*, Vol. 120, Pp. 12025-12038.
- [28]. Walling, D.E., He, Q., Quine, T.A., Irvine, R.J. (2001). "Conversion Of Radionuclide Inventories Measured In One Unit To Another Unit: A Computer Programme." *Journal Of Environmental Radioactivity*, 53(2), 155-165.
- [29]. Reis, J. F., & Cerdà, A. (2013). Influence Of Slope Angle And Length On Sediment And Water Losses Under Different Rainfall Intensities. *Catena*, 110, 70-77.
- [30]. Wu, J., Sun, Y., Yang, X., & Yu, H. (2018). Soil Erosion Processes On Steep Slopes In Subtropical China: A Review. *Journal Of Soils And Sediments*, 18(9), 2929-2941.
- [31]. Bationo, A., Kihara, J., Vanlauwe, B., Waswa, B., Kimetu, J., & Okalebo, J. (2007). Soil Organic Carbon Dynamics, Functions And Management In West African Agro-Ecosystems. *Agricultural Systems*, 94(1), 13-25.
- [32]. Mekuria, W., & Aynekulu, E. (2019). Land Use And Land Cover Effects On Soil Quality In Ethiopia. *Environmental Systems Research*, 8(1), 13.
- [33]. Zhang, X., Hu, Y., Li, H., & Li, C. (2019). Effects Of Conservation Tillage Practices On Soil Carbon Sequestration And Crop Yield In China: A Meta-Analysis. *Journal Of Cleaner Production*, 240, 118204.