

Evaluation of the Outcome of Proximal Humerus Fracture Managed By Internal Locking Plate- A Prospective Study

Dr. T M Jose. *Professor & HOD, Department of Orthopaedics, Jubilee Mission Medical College , Thrissur.*

Dr. Ajith Vincent T. *Junior Resident, Department of Orthopaedics, Jubilee Mission Medical College , Thrissur*

Dr Anil George Paul, *Senior Resident, Department of Orthopaedics, Jubilee Mission Medical College , Thrissur*

Abstract

Background And Objective: Proximal humerus fractures accounts for about 4 to 5% of all fractures. They are the third most common fractures in elderly population after hip and distal radius fractures. Regarding treatment of proximal humerus fractures, controversies still exists whether to do conservative or operative management. Various operative procedures are carried out, recent trend in internal fixation has moved on to locking plates which gives angular stability. The present study is undertaken to evaluate the functional outcome and complications of proximal humerus fractures treated by PHILOS locking plate.

Methods: Prospective study involving Adults(18-60years) with proximal humerus fractures admitted to Jubilee Mission Medical College And Research Institute in the period from January 2015 to December 2015. In this study 30 cases of fractures of proximal humerus were treated by open reduction and internal fixation with PHILOS locking plate and followed up for six months. Patients were followed up in outpatient department at first, second, third and sixth months Serial radiographs were taken for assessing complications and the functional outcomes were assessed using Constant and Murley score.

Results And Discussion: In our series, male female incidence were equal with majority of patients sustaining fracture following road traffic accident. Average age of patients were 43 ± 11.65 and most common fracture pattern observed in these patients were 3 part fracture (33.3%). 66% of the fractures united within three months and 94% of the fracture united within six months. Average constant score of 65.13 was achieved. Overall complication rates were 20% with one case of osteonecrosis of humeral head. 10% of the patient needed a second unplanned surgery. Fracture pattern and complications had significant impact on the functional outcome of the patient.

Conclusion: Internal fixation with PHILOS plate is reliable operative option for displaced proximal humerus fracture which provides angular stable fixation, especially in a osteoporotic bone. High rate of complications depends both on implant design and operative technique. Therefore, attention on technical aspects of applying them would help optimization of the results.

Keywords: Proximal humerus fractures, PHILOS plating, complications, prospective study

I. Introduction

Proximal humeral fractures account for approximately 5% of all fractures. Within the last three decades, the age-adjusted incidence of proximal humeral fractures increased by 15% per year. Increased incidence of proximal humeral fractures is associated with more complications. Up to 80% of proximal humeral fractures can be treated nonoperatively, resulting in satisfactory results.^[3] However, different techniques have been described for fixation of comminuted and displaced proximal humeral fractures, including sutures, cerclage wires, K-wires, screws and plates, intramedullary devices, and shoulder arthroplasty. The complication rate can be as high as 50% or higher.^[8,9] Several complications have been reported, such as cut-out or back-out of the screws and plates, nonunion, avascular necrosis, nail migration, rotator cuff impairment and impingement syndrome.^[10,11] Even shoulder arthroplasty in proximal humeral fractures may yield functionally poor results.^[12] In order to decrease the incidence of complications, particularly fixation failure and loss of stability, and to improve stability and enable early post-operative mobilization, new plating techniques such as the Proximal Humeral Internal Locking System (PHILOS, Synthes, Solothurn, Switzerland) have been developed.^[13] Since there is a high correlation between the holding capacity of screws and regional bone morphology (e.g. cortical thickness and bone mineral density),^[14]

Osteoporotic bone is implicated in the occurrence of complications in proximal humeral fractures. The aim of the present study was to evaluate the clinical results of PHILOS plate fixation in proximal humeral fractures and to analyze potential complications.

I. Objective

The Functional Outcome and Complications Of Proximal Humerus Fracture Treated With PHILOS Plate

II. Methodology

First thirty patients who attended Jubilee Mission Medical College with proximal humerus fracture, satisfying inclusion criteria, during the period of January 2015 to December 2015 were selected for the study. The patient admitted with proximal humerus fracture were examined according to the protocol.

Inclusion Criteria

- Age group : 20 -60 years of age
- Closed fracture
- 2 part, 3 part and 4 part fracture
- fracture dislocation

Exclusion Criteria

- Undisplaced fracture
- Open fracture
- fracture more than 3 weeks
- severely comminuted fracture
- unfit for surgery
- Head injury
- Neurovascular injury
- Previous shoulder pathology

General information like name, age, sex, occupation and address were noted. Then detailed history regarding the mode of injury was elicited. History of the past illnesses were also elicited in detail. Clinical examination of the injured shoulder was done and neurovascular injury was ruled out. Other associated injuries were noted. Radiological evaluation was done and fracture of the proximal humerus was classified according to Neer classification. Blood investigation like haemoglobin, total count, differential count, ESR, random blood sugar, and ECG were done. Screening test were also done for all patient before surgery. All patients underwent a pre-anesthetic check-up by the anesthetist and were operated as early as possible if general conditions were stable.

Preoperative preparations

- patients were kept nil per oral for 6 hours prior to surgery
- An informed consent for surgery was taken
- Systemic antibiotic (ceftriaxone sulbactam combination) was given one hour prior to surgery.

Operative Techniques

The choice of anesthesia was decided by the anesthetist and they have given both general anesthesia as well as regional block in combination. This helped to decrease the immediate post operative pain.

Patient position and drape

Patients placed in supine position on operating table with wedge a sand bag under the spine and medial border of scapula to push the affected side forward while allowing the arm to fall backward. The entire shoulder girdle prepared including the proximal part of the arm, so that incision could be extended. Drape the arm free, because it will have to be moved during the approach.

Surgical approach

Deltopectoral approach- Incision starts just above the coracoid process, which is palpated in deepest point in the clavicular concavity distally towards acromioclavicular joint. An 8 to 10 cm incision started from

just above coracoid process advanced following the line of deltapectoral groove. The intravenous plane is between the deltoid muscle which is supplied by axillary nerve and the pectoralis major muscle, which is supplied by the medial and lateral pectoral nerves. Retract pectoralis major medially and deltoid laterally, splitting the two muscle apart. The vein is retracted either medially or laterally. The short head of biceps and the coracobrachialis must be displaced medially before access can be gained to anterior aspect of shoulder joint. Beneath the tendons lie the transversely running fibers of subscapularis muscle. Apply external rotation to the arm to stretch the subscapularis, bringing the muscle belly into wound and making its superior and inferior borders easier to define. Pass a blunt instrument between the capsule and the subscapularis, then divide the subscapularis in from insertion onto the lesser trochanter of humerus. Incise the capsule longitudinally to enter the joint wherever the selected repair must be performed.

Procedure

Through delto-pectoral approach, the fracture was exposed and reduced with minimal soft tissue dissection. Briefly, the anatomical relationship between humeral head and greater tuberosity was reduced and fixed temporarily with K wires. In case of obvious rotation or displacement of the humeral head, a joystick technique was used. Then the shaft fragment was reduced by abduction, traction and rotation of the arm. Reduction was checked under image intensifier. Definitive fixation with locking plate was done with plate positioned lateral to bicipital groove sparing tendon of long head of biceps and 1cm distal to greater tuberosity. The screws were chosen according to preoperative planning, and all the four head screws were supposed to be inserted to the head fragment. The inferior screws supporting the humeral head were considered critical. Proximal locking screws were inserted to hold the humeral head, which are multi directional screws with the tips of the screws staying 5–10 mm away from the articular surface. All proximal locking screws were placed in a unicortical fashion through an external guide and confirmed to be within the humeral head with intra operative fluoroscopy. AP (internal and external rotation) view used to visualize screw placement. The distal shaft screws were placed bicortically. A minimum of three bicortical screws were used. Fluoroscopic images were taken to confirm satisfactory fracture reduction, plate positioning and proper length of screws in the humeral head. In case of severe comminution or instability, the rotator cuff, the greater tuberosity, and the lesser tuberosity were fastened to the plate using non-absorbable sutures. Range of motion of shoulder was checked on the table for impingement. Wound was closed under negative suction, which was removed after 48 hours.

Postoperative management

All patients are immobilized in arm pouch with cuff and collar sling. Appropriate antibiotics and analgesics were used. Immediate post operative radiographs were taken to determine the bone alignment and maintenance of reduction. Sutures removed by 10th day. Passive range of motion and pendulum exercises are begun immediately depending on pain. All patients were followed up at 4 weeks, 8 weeks, 12 weeks and at 6 months.

Exercise regime post PHILOS plating

From post operative day one onwards patients are started on pendulum exercises with their arm pouch until the first follow-up visit in the op. During the first visit arm pouch was removed and started on active assisted external rotational to neutral. From two months onwards patients were allowed to have full range of movements.

Functional Outcome

Final results are assessed using the Constant and Murley scoring system

Fracture Union:

At first patients are clinically evaluated for fracture union. Then plain radiographs of shoulder are taken in anteroposterior and axillary views to decide on fracture union. If three out of four cortices shows bridging, fracture is considered as united. This is done on first, second, third and sixth month of follow up.

III. Results

Seventy three patients with proximal humerus fracture were treated in the institution during the period of January 2015 to December 2015. Forty three patient were excluded on the basis of exclusion criteria in which one was a open two part fracture following a sharp cut injury and one fracture associated with subdural hemorrhage. Most of the excluded patients were above the age of sixty who were injured after a trivial fall. Therefore, first thirty patients with inclusion criteria were treated with open reduction and internal fixation with PHILOS plate. The mean age of all the patient in this study was found to be 43 ± 11.65 . The proximal humeral fracture was following a road traffic accident in 16(53%) patients, a fall from standing height in 12(40%) patients

and seizure in 2(6.66%)patients.17 patients had injury to their dominant side. Dominant and non-dominant sides were equally affected. Gender distribution males were more involved.

Out of the thirty patients, 26% sustained two part fracture,33.3% had three part fracture,16% had four part fracture and 23.3% had fracture dislocation. The most common fracture type was 3-part fractures. The average hospital stay for patients were 8.2 days. At the one month, two month, three month and six month follow-up examinations, twenty nine out of thirty patients came for follow up. One of the patients in the study was lost in follow up.

During the follow up the Constant score improved significantly from the first month to the sixth month.The mean Constant score at 6th month of follow-up was 65.13±15.17. There was no statistically significant difference in the functional outcome of different age groups.

Table No: Age specific constant score

Age	Constant score				Total
	Excellent	Fair	Good	poor	
21-30	3	2	2	0	7
31-40	1	4	0	1	6
41-50	0	2	3	2	7
51-60	5	2	2	1	10
Total	9	10	7	4	30

Pearson Chi-Square value is 11.7; p value = 0.227

There was statistically significant association between fracture type and functional outcome.

Table no: relationship between types of fracture and average constant score

	No. of cases	Percentage	Average constant score	Inference
2 part	8	26%	9	excellent
3 part	10	33.3%	16	Fair
4 part	5	16%	23	good
Dislocation	7	23%	28	good
Total	30	100%	76	

Out of the 30 patients, 8patients with 2 part fracture had an average constant score of 9 which was excellent. 10 patients had 3part fracture with an average constant score of 16 which was fair in outcome. 4part and fracture dislocation attained a lower constant score with good outcome.

This correlation indicated that, more complex the fracture, the functionoutcome was affected, and simple fracture had better functional outcome. The number of united fracture increased from 66 % at third month to 96% at the sixth month of follow up. There is astatisticallysignificant difference in the union rates of different fracture types.

Neer classification	Union in Weeks				Total
	8	12	24	>24	
1	2	3	3	0	8
2	3	7	0	0	10
3	0	5	0	0	5
4	1	1	3	2	7
Total	6	16	6	2	30

Pearson Chi-Square value is 19.27; p value = 0.023

Complications

Six complications(20%) were encountered in Thirty study patients.

Complication	Frequency	Percent
Nil	24	80.0
Yes	6	20.0
Total	30	100.0

Two patients had deep infection which was evident by 2nd month of follow up and required intravenous antibiotic and eventually implant was removed. Post surgery, one patient sustained brachial plexus injury. One patient had shoulder joint subluxation in the third month. One patient had avascular necrosis of humeral head which was evident in the 6th month of follow-up. Except for one patient with subacromial impingement because of cranially placed Locking plate, there was no other implant related complications. No patient had primary screw perforation of humeral head which was unrecognized intraoperatively. In total, three(10%) patient had under gone second unplanned surgery with in six months after first operation. Patient without any complications attained statistically significant higher Constant score than patients having any one of the complications.

Complication	Functional outcome				Total
	Excellent	Fair	Good	poor	
0	9	9	5	1	24
1	0	1	2	3	6
Total	9	10	7	4	30

Pearson Chi-Square; 10.72 p value is 0.013

Patients with complications did not show statistically significant difference in union of fracture.

Complication	Fracture union at 6 th month		Total
	nonunion	Union	
No	0	24	24
Yes	2	4	6
Total	1	28	30

Fisher's Exact Test; p value is 0.0759

Complications and duration for surgery did not show any association. But it showed 40% positive correlation.

complication	Time in minutes					Total
	<90	90-120	121-150	150-180	>180	
No	1	7	11	2	3	24
Yes	0	1	0	2	3	6
Total	1	8	11	4	6	30

Pearson Chi-Square 8.9; p value 0.063

Spearman Correlation + 0.40

IV. Discussion

Proximal humeral fractures represent an increasing challenge for the health-care system because of the increasing proportion of elderly individuals in the population. As estimated by a Finnish study group, the number of fractures may triple by the year 2030¹⁵. The majority of patients with these fractures are more than sixty years old, and most of the proximal humeral fractures in this population are related to osteoporosis¹⁶. Nevertheless, stable reduction is essential for healing of the fracture and for achieving early functional recovery of the shoulder. In patients with osteoporotic bone and/or comminuted fractures, operative stabilization is challenging and the treatment of displaced and unstable fractures remains controversial.

Various techniques have been used to stabilize fractures of the proximal part of the humerus, including intramedullary nails, plate-and-screw osteosynthesis, tension band wiring, percutaneous pin fixation, and hemiarthroplasty^{17,18,19}. Successful outcomes after plate osteosynthesis of proximal humeral fractures have been reported^{19,20,21}. Open reduction and internal fixation of proximal humeral fractures with non-locking plates and screws has been shown to provide the strongest fixation in non-osteoporotic bone¹⁷. In that study, the average age of the patients at the time of the injury was forty- eight years. As the stability of the osteosynthesis with non- locking plates and screws relies on the friction between the plate and bone, the effectiveness of traditional plate-and-screw fixation decreases with bone quality. Additionally, complications such as screw loosening resulting from the insufficient purchase of screws in osteoporotic bone lead to high failure rates, especially among patients with three and four-part fractures. Kristiansen and Christensen¹⁴ reported satisfactory or excellent results in only nine of twenty patients who had fixation of a proximal humeral fracture with a T-

buttress plate, and there was a high rate of fixation failure. The authors stated that the results for all four-part fractures were poor and recommended primary treatment with a prosthesis.

New techniques involving the use of plates and screws with angular stability have been introduced in order to avoid these complications. The Locking Proximal Humerus Plate was designed to maintain a stable fracture reduction even in osteoporotic bone. Advantages of the Locking Proximal Humerus Plate include gentle fracture reduction with use of indirect maneuvers, a high resistance to avulsion even in patients with poor bone stock because of the combination of fixed-angle screw-plate locking and three-dimensional placement of screws in the humeral head, and the possibility of early exercise and a short period of immobilization because of the high initial stability achieved¹⁵.

The mean age of patients with proximal humerus fracture in our study was 43±11.65 which is not comparable with other studies done previously. The mean age of patients in most of the other studies were more than sixty since this is a fracture of elderly.^{20,21,22,23} We included only patients below the age of sixty because we wanted to avoid skewing in the measured results due to osteoporosis.

In our study there was male preponderance, while most of the other studies had female preponderance.^{20,21,22,23} This is due to the fact that female are predisposed to osteoporosis whom we did not include in our study. One study concluded that female patients were significantly older than male patients⁸⁴ while we could not find any relations in gender age and its incidences. Patients sustaining fracture following high energy trauma were higher compared to other studies in which most of the fracture were following low energy trauma like fall from standing height.^{20,21,22,23} 53% of our patients had fracture following road traffic accident while its only 13 to 25% in other studies. Variation may be due to the exclusion of patients above the age of sixty. Two patients had fractures following seizure episodes which was not reported in any other studies. We could not find any significant difference between the incidence of fracture in dominant and non-dominant sides.

According to Neer classification, 26% two part, 33.3% three part, 16% four part fractures and 23.3% fracture dislocation was found. Study by Felix Brunner et al had similar incidences except for fracture dislocation which was 5% in their study. Most of the patients in our study had a high energy trauma which might be the reason for higher incidence of fracture dislocation compared to other studies. The most common fracture type was 3-part fractures which was consistent with other studies.²³

The mean time between injury and surgery was 2.76 days and did not have any significant influence over the functional outcome of the patients. This was consistent with other studies. Average interval between fracture and surgery was 2.9 days in Felix Burnner et al study. The mean time taken for ORIF was 150.5 minutes. This longer surgery time compared to other studies may be due to the higher incidence of fracture dislocation in our study. No statistically significant association was found between surgery time and complications.

Although our study population consisted patients with mostly complex fracture patterns, the incidence of secondary displacement, pseudarthrosis, or osteonecrosis complications was relatively low compared with previously performed studies concerning the treatment of proximal humerus fractures. 66% of the fracture united within 3 months and 94% of the fracture united within 6 months. We observed six complications(20%) in thirty patients during the six months of followup. There is very high variation between the studies in terms of incidence of complications. Felix brunner et al and Plecko et al²⁴ detected 33% and 34% complications respectively in patients with PHILOS plate. One study mentioned that patients over the age of sixty were at a significant 1.9 times higher incidence of experiencing any complications compared to young patients.²⁵ This could be the reason for the lower incidence of complications in our study. In our study, there was a case of subacromial impingement which was directly related to the initial surgical procedure itself. The patients score improved once the implant was removed.

The main challenge in the operative treatment of proximal humeral fractures is to achieve effective stabilization of an adequately reduced fracture in order to maximize the functional outcome. If an adequate reduction is not achieved and medial buttressing is insufficient, especially with a varus malreduction, secondary loss of reduction and subsequent screw perforation or plate breakage is possible. The locking of the screws onto the plate prevents the screws from backing out. Therefore, if the fracture collapses, the screws may penetrate the articular surface. This penetration may be more likely if the screws are placed very close to the articular surface or if the articular surface was penetrated during drilling.

The most common complications reported in most of the other studies were screw perforation either primary or secondary.^{21,22} The working groups of Kettler et al²⁶ and Charalambous et al²⁷ detected 24/176 and 2/17 primary screw perforations, respectively, in their series of PHILOS plate patients. Most of this patients were over the age of sixty which we did not include. Fortunately we did not have a case of primary perforation because of the meticulous intraoperative screening using c-arm. We also did not have a case of secondary perforation of humeral head with screws, which is usually seen in patients with poor bone stock or following varus malreduction during the initial surgery.

Avascular necrosis is the one of the most feared complications following open reduction internal fixation of proximal humerus fractures. AVN can develop as long as five years after injury.²⁸ This phenomenon is supported by the observation that studies with longer follow up usually also report higher rates of AVN. In our study we had one patient with avascular necrosis which was diagnosed at sixth month followup. The incidence of AVN varies from 0-68% throughout the literature^{30,31}. The fracture type itself, the length of the dorsomedial metaphyseal extension, and the integrity of the medial hinge were found to be the most relevant predictors of humeral head necrosis. This is supported by the a study by Jost et al³². presenting a AVN rate of 68% including only three and four part fractures. But in our study we could not find a relation between fracture type and occurrence of AVN. The Constant score for AVN patient was 66. Wijnman et al⁷⁸ reported that few patients with radiographic evidence of AVN had a good or excellent Constant score compared to those with AVN

V. Conclusion

The present study was done to evaluate functional outcome following surgical management of displaced proximal humerus fracture by PHILOS locking plate. In our study of 30 patients below the age of sixty with displaced proximal humerus fractures treated with proximal humeral internal locking system plating, we conclude that the functional outcome is much better in patients with 2 part and 3 part fractures, and not good in complex fractures and fracture dislocations. The advantage of this fixation is it allows early postoperative mobilization of the affected shoulder and better functional outcome of the affected shoulder, as compared to conservative management, where patients affected shoulder is immobilized for long periods³⁴. Among the 30 patients 6(20%) had complications, and 3 had unplanned second surgery. Proximal Humeral Internal Locking system (PHILOS); In this system, locking of the threaded heads of the screws in the plate itself provides for a construct with angular and axial stability, eliminating the possibility of screw toggling (windscreen wiper effect), or sliding of the screws in the plate holes. Coupled with a divergent or convergent screw orientation, this makes for much improved resistance to pull out and failure of fixation³⁵. Also, whereas conventional plating systems depend on compression between the plate undersurface and bone for stability, this is not the case for the PHILOS. This lessens the chance of stripping the thread in osteoporotic bone, as the plate/bone interface is not loaded along the screw axis. This also allows for a more biological fixation as the underlying periosteum and blood supply to the fractured regions are much less compressed.

To conclude, PHILOS locking plate is a preferential implant in proximal humeral fractures due to angular stability, particularly in comminuted fractures in younger patients, thus allowing early mobilization.

Bibliography

- [1]. Gerald R. Williams and Kirk L. Wong, 2000: "Two-part and three part fractures- Management of proximal and distal humerus fracture". *Orthop Clin North Am*, January 31 (1) : 1-21.
- [2]. Lind T, Kronerker, Jensen J. : The epidemiology of fractures of proximal humerus. *Arch Orthop Trauma Surg*, 1989; 108: 285.
- [3]. Louis U., Bigliani, Chapter 9 The shoulder, Vol-1 ed. Charles Rockwood, Frederick A. Fractures of proximal humerus. In Rockwood CA, Matsen. Philadelphia: W.B. Saunders, 1990: p278-334.
- [4]. Scott E. Powell, Robert W. Chandler. Fractures of the proximal humerus. Chapter- 11, In: Text book of Operative techniques in upper extremity sports injuries. Ed.Frank W. Jobe, Mosby, 1995: p.313-340.
- [5]. Zyto K. Non-operative treatment of comminuted fracture of proximal humerus in elderly patients. *Injury*, 1998; 29: 349-52.
- [6]. Zyto K, Wallace WA, Frostick SP, Preston BJ. Outcome after hemiarthroplasty for three and four part fracture of the proximal humerus. *J Shoulder Elbow Surg*, 1998; 7: 85-9
- [7]. Wijnman A.J, Roolker W, Patt T W et al , open reduction and internal fixation of three and four part fractures of proximal humerus. *Scientific Article* November 01, 2002 106 8. Sarah L.Edwards, Nicole A.wilson, Li-gun Zhang et al Two part surgical neck fractures of proximal part of humerus: Biomechanical Evaluation of Two Fixation. *The journal of bone and joint surgery* 2006 88 :2258-64
- [8]. Gerber C, Worner CM, Vienne P. "Internal fixation of complex fractures of the proximal humerus". *J Bone Joint Surg (Br)*, 2004 Aug; 86(60):P 848-55.
- [9]. Neer CS. Displaced proximal humeral fracture : Part I: Classification and evaluation. *J Bone Joint Surg (Am)* 1970; 52-A: 1077-89.
- [10]. Neer CS. "Displaced proximal humeral fractures. Part II. Treatment of three part and four part displacement". *J Bone Joint Surg*, 1970; 52A: 1090-1103.
- [11]. Paavolainen P, Bjorkenheim JM, Slati P, and Paukku P. "Operative treatment of severe proximal humeral fractures". *Acta Orthop Scand*, 1983; 54: 374-379. 13. Siebler G, Kuner EH , "Late results following the surgical treatment of proximal humerus fractures in adults". *Unfall chirurgie* 1985; 11(3): 119-127.
- [12]. Kristiansen B and Christensen SW : Plate fixation of proximal humeral fractures, *Acta Orthop Scand*, 1986; 57: 320-323,
- [13]. Palvanen M, Kannus P, Niemi S, Parkkari J. Update in the epidemiology of proximal humeral fractures. *Clinical orthopaedics and related research*. 2006 Jan 1;442:87-92.
- [14]. Jones G, Nguyen T, Kelly PJ, Gilbert C, Eisman JA. Symptomatic fracture incidence in elderly men and women: the Dubbo Osteoporosis Epidemiology Study (DOES). *Osteoporosis International*. 1994 Sep 1;4(5):277-82.
- [15]. Koukakis A, Apostolou CD, Taneja T, Korres DS, Amini A. Fixation of proximal humerus fractures using the PHILOS plate: early experience. *Clinical orthopaedics and related research*. 2006 Jan 1;442:115-20.
- [16]. Resch H, Povacz PA, Fröhlich R, Wambacher MA. Percutaneous fixation of three-and four-part fractures of the proximal humerus. *Bone & Joint Journal*. 1997 Mar 1;79(2):295-300.

- [17]. Wijgman AJ, Roolker W, Patt TW, Raaymakers EL, Marti RK. Open reduction and internal fixation of three and four-part fractures of the proximal part of the humerus. *J Bone Joint Surg Am*. 2002 Nov 1;84(11):1919-25.
- [18]. Wanner GA, Wanner-Schmid E, Romero J, Hersche O, von Smekal A, Trentz O, Ertel W. Internal fixation of displaced proximal humeral fractures with two one-third tubular plates. *Journal of Trauma and Acute Care Surgery*. 2003 Mar 1;54(3):536-44.
- [19]. Hessmann M, Baumgaertel F, Gehling H, Klingelhoefter I, Gotzen L. Plate fixation of proximal humeral fractures with indirect reduction: surgical technique and results utilizing three shoulder scores. *Injury*. 1999 Sep 30;30(7):453-62.
- [20]. Südkamp N, Bayer J, Hepp P, Voigt C, Oestern H, Kääh M, Luo C, Plecko M, Wendt K, Köstler W, Konrad G. Open reduction and internal fixation of proximal humeral fractures with use of the locking proximal humerus plate. *J Bone Joint Surg Am*. 2009 Jun 1;91(6):1320-8.
- [21]. Sproul RC, Iyengar JJ, Devcic Z, Feeley BT. A systematic review of locking plate fixation of proximal humerus fractures. *Injury*. 2011 Apr 30;42(4):408-13.
- [22]. Thanasis C, Kontakis G, Angoules A, Limb D, Giannoudis P. Treatment of proximal humerus fractures with locking plates: a systematic review. *Journal of Shoulder and Elbow Surgery*. 2009 Dec 31;18(6):837-44.
- [23]. Björkenheim JM, Pajarinen J, Savolainen V. Internal fixation of proximal humeral fractures with a locking compression plate A retrospective evaluation of 72 patients followed for a minimum of 1 year. *Acta orthopaedica Scandinavica*. 2004 Jan 1;75(6):741-5.
- [24]. Plecko M, Kraus A. Internal fixation of proximal humerus fractures using the locking proximal humerus plate. *Operative Orthopädie und Traumatologie*. 2005 Feb;17(1):25-50.
- [25]. Brunner F, Sommer C, Bahrs C, Heuwinkel R, Hafner C, Rillmann P, Kohut G, Ekelund A, Müller M, Audigé L, Babst R. Open reduction and internal fixation of proximal humerus fractures using a proximal humeral locked plate: a prospective multicenter analysis. *Journal of orthopaedic trauma*. 2009 Mar 1;23(3):163-72.
- [26]. Kettler M, Biberthaler P, Braunstein V, Zeiler C, Kroetz M, Mutschler W. [Treatment of proximal humeral fractures with the PHILOS angular stable plate. Presentation of 225 cases of dislocated fractures]. *Der Unfallchirurg*. 2006 Dec;109(12):1032-40.
- [27]. Charalambous CP, Siddique I, Valluripalli K, Kovacevic M, Panose P, Srinivasan M, Marynissen H. Proximal humeral internal locking system (PHILOS) for the treatment of proximal humeral fractures. *Archives of orthopaedic and trauma surgery*. 2007 Apr 1;127(3):205-10.
- [28]. Patel S, Colaco HB, Elvey ME, Lee MH. Post-traumatic osteonecrosis of the proximal humerus. *Injury*. 2015 Oct 31;46(10):1878-84.
- [29]. Clavert P, Adam P, Bevort A, Bonnomet F, Kempf JF. Pitfalls and complications with locking plate for proximal humerus fracture. *Journal of Shoulder and Elbow Surgery*. 2010 Jun 30;19(4):489-94.
- [30]. Grawe B, Le T, Lee T, Wyrick J. Open reduction and internal fixation (ORIF) of complex 3-and 4-part fractures of the proximal humerus: does age really matter?. *Geriatric orthopaedic surgery & rehabilitation*. 2011 Dec 19;2151458511430662.
- [31]. Acklin YP, Stoffel K, Sommer C. A prospective analysis of the functional and radiological outcomes of minimally invasive plating in proximal humerus fractures. *Injury*. 2013 Apr 30;44(4):456-60.
- [32]. Jost B, Spross C, Grehn H, Gerber C. Locking plate fixation of fractures of the proximal humerus: analysis of complications, revision strategies and outcome. *Journal of shoulder and elbow surgery*. 2013 Apr 30;22(4):542-9.
- [33]. Sharafeldin KN, Quinlan JF, Corrigan J, Kelly IP. Functional follow-up of locking plate fixation of fractures of the proximal humerus. *European Journal of Orthopaedic Surgery & Traumatology*. 2008 Feb 1;18(2):87-92.