

Preemptive Diclofenac in Men Undergoing Double J Stent Removal- A Randomized Double-Blinded Study

Balaji A.R ¹, Chandramurali R ²

¹Professor And Head Of The Department, Department Of Urology, Stanley Medical College, Chennai, Tamil Nadu

²Post Graduate, Department Of Urology, Stanley Medical College, Chennai, Tamil Nadu

Abstract:

Introduction: Double J (DJ) stent removal requires local anesthesia and is done using rigid cystoscope. Patients experience significant pain during and after the procedure. We assessed the efficacy of preemptive oral diclofenac in analgesia for patients undergoing DJ stent removal by rigid cystoscopy compared to placebo.

Methods: Only male patients undergoing DJ stent removal under local anesthesia between March 2015 and July 2016 were enrolled. Patients were randomized to receive 75 mg oral diclofenac (Group A) or placebo (Group B) 1 hour before procedure by randomization. Intraurethral 2% lignocaine gel (25 ml) was used in both groups. Pain during rigid cystoscopy, pain at the first void, and at 1 day after cystoscopy was assessed using visual analog scale (VAS) (0–100). Adverse reactions to diclofenac and episodes of acute urinary retention, if any, were assessed.

Results: Totally 121 males (Group A [n = 62]; Group B [n = 59]) underwent stent removal. The median (Interquartile range) VAS during the procedure in Group A was 30 (30) and Group B was 60 (30) (P < 0.001), at first void was 30 (30) and 70 (30) (P < 0.001) and at 24 h postoperatively was 20 (20) and 40 (20) (P < 0.001). The incidence of side effects was comparable in the two groups (P > 0.05).

Conclusions: A single oral dose of diclofenac 1 h before DJ stent removal using rigid cystoscope under intraurethral lignocaine anesthesia decreases pain significantly during and upto one day after the procedure, with minimal side effects.

Keywords: Analgesia, Double J stent, Diclofenac

I. Introduction

Double J (DJ) stents are removed under local anesthesia using a rigid cystoscope. Patients experience significant pain during and after the procedure for few days. Lignocaine gel during this procedure is not very effective.^{[1],[2],[3],[4],[5]} Numerous reports exist on the effect of local anesthesia against cystoscopy-associated pain, but so far there has been no study on the preemptive analgesic effect of nonsteroidal anti-inflammatory drugs (NSAIDs).^{[1],[2]} Hence, our study was done to assess the effect of preemptive oral diclofenac for analgesia in patients undergoing DJ stent removal by rigid cystoscopy when compared to placebo and to assess its safety.

Methods Trial design

The study was conducted from March 2015 to August 2016 in the Department of Urology, Government Stanley medical college, Chennai in 121 patients undergoing DJ stent removal using rigid cystoscope.

Study design: double-blind, randomized, placebo-controlled trial

Inclusion criteria

All random male patients aged more than 18 years, undergoing DJ stent removal under local anesthesia (2% lignocaine gel), were eligible for the study after providing a written informed consent.

Exclusion criteria

The exclusion criteria were age < 18 years, inability to understand visual analog scale (VAS) for pain, lower urinary tract symptoms (LUTS) before initial surgery, prostatitis or urethral stricture during cystoscopy, renal failure (serum creatinine > 1.5 mg/dl), concomitant usage of nephrotoxic drugs, history suggestive of or known case of peptic ulcer disease or upper gastrointestinal bleed or upper abdominal surgeries. Patients were randomized to receive oral diclofenac 75 mg or placebo by randomization. The patient and the investigator who assessed the outcome (pain) were also blinded.

Intervention

All patients were administered the drug or placebo 1 h before the procedure. A 20 Fr cystoscope was used for DJ stent removal in all patients. 25 ml of 2% lignocaine gel was injected intraurethrally and was allowed to act for 10 minutes. Cystourethroscopy was performed and DJ stent was removed with a DJ stent

removal forceps. Postoperative pain score was evaluated with the 100-point VAS.^{[6],[7]} VAS was assessed during procedure, first void, and 24 h after DJ stent removal. The presence of nausea, vomiting, epigastric pain, and urinary retention was recorded.

II. Statistical analysis

Statistical analysis was performed using SPSS version 20.Independent samples t-test or Mann–Whitney U-test as appropriate was used based on the normality of the distribution. Chi-square/Fisher's exact test was used for categorical variables. The P< 0.05 was considered statistically significant.

III. Results

The baseline patient characteristics such as age, side, weight, duration of stent, and operative time were almost similar[Table 1].

Patient characteristics	Groups		P
	A (n=62)	B (n=59)	
Age, mean±SD (years)	47.5±9.76	46.05±9.12	0.22
Right:left (n)	41:21	30:29	0.47
Duration of stent, mean±SD (days)	20.93±5.36	21.36±5.45	0.87
Weight, mean±SD (kg)	51.35±10.12	51.75±10.49	0.18
Duration of procedure, mean±SD (min)	2.30±1.20	2.50±1.31	0.63

SD=Standard deviation

Pain

The median pain score was lower in the Group A during, at first void, and 1 day after DJ stent removal. Statistically significant difference was observed between Groups A and B [Table 2].

Timing	Visual analog scale		P
	A (n=62)	B (n=59)	
Pain during procedure, median (IQR)	30 (30)	60 (30)	<0.001
Pain during first void, median (IQR)	30 (30)	70 (30)	<0.001
Pain after 24 h, median (IQR)	20 (20)	40 (20)	<0.001

IQR=Interquartile range

Adverse reactions

Epigastric pain, nausea and vomiting were slightly higher in the patients receiving diclofenac, but not statistically significant. No patient developed acute urinary retention. [Table 3].

Adverse reactions	Number of patients, n (%)		P
	A (n=62)	B (n=59)	
Epigastric pain	10 (16)	6 (10)	0.08
Nausea and vomiting	8 (13)	5 (8.5)	0.06

IV. Discussion

This study was performed to assess if diclofenac is effective in reducing pain during DJ stent removal using rigid cystoscopy. Preemptive analgesia before cystoscopy under local anesthesia should be considered as the standard of care to reduce pain experienced by the patient. Lignocaine gel has little analgesia only.^{[2], [3], [4]} So far there has been no objective evaluation of the preemptive analgesic effect of NSAID in rigid cystoscopy, pain score during cystoscopy was maximum, followed by the first and second urination after cystoscopy, injection of gel, and urination the day after cystoscopy.^[2] We found that oral diclofenac was effective in reducing pain during stent removal and up to 24 h after the procedure in our patients. We included only male patients to avoid the confounding effect of urethral length in assessing the efficacy of diclofenac.

Surgical trauma induces cyclooxygenase-2 (COX-2) enzyme causing release of prostaglandins, which sensitize the peripheral nociceptors and produce localized primary hyperalgesia. This resultant central sensitization produces pain hypersensitivity in the surrounding uninjured tissue.^[8] After nociception, central sensitization induces c-Fos expressed in spinal dorsal neurons which augments the nociceptive sensitivity. Wind-up is the mechanism of central sensitization, mainly by the activation of N-methyl-D-aspartate (NMDA) and neurokinin receptors. Preoperative NSAIDs before tissue trauma set in prevent the development of hyperalgesia.^[8]

NSAIDs, by inhibiting the early production of prostanoids before surgical trauma more effectively, prevent the development of both peripheral and central sensitization and hyperalgesia. They also block the nociceptive impulses, increase the threshold of nociceptive neurons by opioids, block the wind-up using NMDA receptor antagonists, and suppress local inflammation. It is possible to produce a painless postsurgical state with NSAIDs. Diclofenac is a nonspecific COX inhibitor. Since patients are preemptively treated with diclofenac, inflammation incited by the stent removal procedure is lesser, and consequently, pain incited is lesser. In our study, pain assessment by VAS was done up to 24 h.

V. Conclusion

A single preprocedural oral dose of diclofenac administered 1 h before DJ stent removal done by rigid cystoscope under intraurethral lignocaine anesthesia decreases pain significantly during and up to 24 h postprocedure with minimal side effects and can be safely prescribed for office endo-urological procedures.

References:

- [1]. Kobayashi T, Nishizawa K, Mitsumori K, Ogura K. Instillation of anesthetic gel is no longer necessary in the era of flexible cystoscopy: A crossover study. *J Endourol* 2004;18:483-6.
- [2]. Chen YT, Hsiao PJ, Wong WY, Wang CC, Yang SS, Hsieh CH. Randomized double-blind comparison of lidocaine gel and plain lubricating gel in relieving pain during flexible cystoscopy. *J Endourol* 2005;19:163-6.
- [3]. McFarlane N, Denstedt J, Ganapathy S, Razvi H. Randomized trial of 10 mL and 20 mL of 2% intraurethral lidocaine gel and placebo in men undergoing flexible cystoscopy. *J Endourol* 2001;15:541-4.
- [4]. Herr HW, Schneider M. Outpatient flexible cystoscopy in men: A randomized study of patient tolerance. *J Urol* 2001;165(6 Pt 1):1971-2.
- [5]. Komiya A, Endo T, Kobayashi M, Kim W, Araki K, Naya Y, et al. Oral analgesia by non-steroidal anti-inflammatory drug zaltoprofen to manage cystoscopy-related pain: A prospective study. *Int J Urol* 2009;16:874-80.
- [6]. Aubrun F, Langeron O, Quesnel C, Coriat P, Riou B. Relationships between measurement of pain using visual analog score and morphine requirements during postoperative intravenous morphine titration. *Anesthesiology* 2003;98:1415-21.
- [7]. Jensen MP, Chen C, Brugger AM. Interpretation of visual analog scale ratings and change scores: A reanalysis of two clinical trials of postoperative pain. *J Pain* 2003;4:407-14.
- [8]. Aida S, Baba H, Yamakura T, Taga K, Fukuda S, Shimoji K. The effectiveness of preemptive analgesia varies according to the type of surgery: A randomized, double-blind study. *Anesth Analg* 1999;89:711-6.
- [9]. Ong CK, Lirk P, Seymour RA, Jenkins BJ. The efficacy of preemptive analgesia for acute postoperative pain management: A meta-analysis. *Anesth Analg* 2005;100:757-73.
- [10]. Reuben SS, Bhopatkar S, Maciolek H, Joshi W, Sklar J. The preemptive analgesic effect of rofecoxib after ambulatory arthroscopic knee surgery. *Anesth Analg* 2002;94:55-9.