

Sponataneous perforation of Meckel's diverticulum: A Case report

Dr.Vineetkumar Mishra¹, Dr.Yogender Kadian², Dr.Nikhil Tayal³,
Dr.Arnab Kadian⁴

^{1,3}(Department of General Surgery,Pt.B.D.SharmaPGIMS,Rohtak/UHSR,India)

²(Department of peadiatric surgery,Pt.B.D.Sharma PGIMS)

Corresponding Author: Dr.Vineetkumar Mishra

Abstract: Meckel's diverticulum is a commonest abnormality of gastrointestinal tract. The incidence of Meckel's diverticulum (MD) varies between 1 and 2% and carries the lifetime risk of 4–6% to become symptomatic. The complications associated with MD include inflammation, perforation, hemorrhage, intussusception, volvulus, intestinal obstruction, and malignant transformation. A 5 yr old male child presented to paediatric emergency with complaints of pain abdomen for 3 days, green colored vomiting and non passage of stool for 2 days and fever for 1 day. A diagnosis of peritonitis was made and decision was made to go for exploratory laparotomy. Operative finding included perforated meckel's diverticulum for which resection and anastomosis were done. Hospital stay was uneventful and patient were discharged. In patients with clinical signs of acute abdomen, Meckel's diverticulum and its potential complications should be kept in mind. Early diagnosis and timely operative intervention must occur in order to provide the best outcome for these patients.

Keywords: Meckel's Diverticulum, Perforation, Peritonitis

Date of Submission: 04-01-2019

Date of acceptance: 19-01-2019

I. Introduction

German anatomist Johann Friedrich Meckel first described the embryological and pathological features in 1809¹. The incidence of Meckel's diverticulum (MD) varies between 1 and 2% and carries the lifetime risk of 4–6% to become symptomatic². Meckel's diverticulum is a commonest abnormality of gastrointestinal tract. It results from patent vitelointestinal duct. It is often found incidentally at the time of abdominal exploration. The complications associated with MD include inflammation, perforation, hemorrhage, intussusception, volvulus, intestinal obstruction, and malignant transformation³. As in acute appendicitis, Meckel's diverticular obstruction results in distal inflammation, necrosis, or even perforation, leading to abscess or peritonitis. Other various pathologies leading to perforation are ulceration of ectopic gastric tissue, ingestion of foreign bodies, Littre's hernia, tumours such as leiomyosarcoma, lymphatic sarcoma, and poorly differentiated stromal tumour^{2,4}

II. Case Report

A 5 yr old male child presented to paediatric emergency with complaints of pain abdomen for 3 days, green colored vomiting and non passage of stool for 2 days and fever for 1 day. There was no history of similar complaints in the past. The patient was a full term vaginal delivery with immediate crying. He had no history of previous hospitalization, jaundice or any surgery. There was no history of breath holding spells. All the developmental milestones were at normal age. There was no history of altered bowel bladder habits. On examination, patient was having a pulse rate of 108/min and respiratory rate of 24/min. On per abdomen examination, distension was present. On palpation, generalized tenderness and guarding was there and bowel sounds were absent.

His blood investigations showed Hb – 10.5gm%, TLC – 9500/mm³, DLC – 60/35/03/02, platelet count – 2.5 lacs/mm³, blood urea – 38mg/dL, random blood sugar – 64mg/dL, Serum sodium – 138mEq/dL, serum potassium – 3.7mEq/dL. On erect x-ray abdomen, there was multiple air fluid levels with no free air under diaphragm. On USG abdomen, there was evidence of free fluid with internal echoes in the pelvis, both paracolic gutter and inter bowel loops which was suggestive of perforation.

A diagnosis of peritonitis was made and decision was made to go for exploratory laparotomy. Operative findings were around 200 cc of purulent fluid in peritoneal cavity with pus flakes over the small gut, liver and spleen. A diverticulum was present in the ileum, around 50 cm from ICJ at the antimesenteric border, which was inflamed and burst at the tip (fig. 1 & 2). The segment of small bowel containing the diverticulum

was resected and anastomosis of the two ends was done. Thorough peritoneal lavage was done. Patient was shifted to paediatric surgery ward after surgery and was kept nil per oral with active ryle's tube aspiration and intravenous fluids and antibiotics, and injection diclofenac for pain relief. Patient's vitals remained stable in the post operative period. Ryle's tube was taken out on 3rd postoperative day and was allowed sips of clear fluid. The oral intake was increased over 3 days as the patient tolerated the feeds. On the 6th post operative day he was fully allowed orally and he was discharged under all satisfactory on the next day.



Figure 1

Figure 2

III. Discussion

Meckel's diverticulum is the most frequent congenital anomaly of the gastrointestinal tract, resulting from the incomplete atrophy of the omphalomesenteric duct. Heterotopic tissues such as gastric, duodenal, colonic ones and rarely pancreatic mucosa can be found in the diverticula, as well as the anatomically normal intestinal mucosa.⁵

Perforation is noted to be an occasional consequence of acute Meckel's diverticulitis, but the exact rate of this has not been reported. The overall lifetime complication rate is approximately 4%. The most common presentation associated with symptomatic Meckel's diverticulum is bleeding, followed by intestinal obstruction, diverticulitis, intussusceptions and neoplasm. However, perforation is very rarely seen and, in a review, was reported as being responsible for 0.5% of symptomatic diverticulum.²

The perforation of a Meckel's diverticulum may mimic acute appendicitis and present as an acute abdomen. The perforation of a Meckel's diverticulum is either caused by; foreign body due to irritation of foreign body and pressure necrosis of the diverticulum wall, or spontaneous perforation due to progressive inflammation of Meckel's diverticulum wall as our case, which produced peritonitis.⁶

A preoperative diagnosis of a complicated MD may be challenging because of the overlapping clinical and imaging features of other acute surgical and inflammatory conditions of the abdomen. A more specific diagnosis, however, will lead to greater recourse to a laparoscopic approach in its treatment.^{6,7}

IV. Conclusion

In patients with clinical signs of acute abdomen, Meckel's diverticulum and its potential complications should be kept in mind. Meckel's diverticulum complications are uncommon and challenge to diagnose. Early diagnosis and timely operative intervention must occur in order to provide the best outcome for these patients. Spontaneous perforated MD often presents as acute abdomen and its preoperative diagnosis is difficult.

References

- [1]. Yorganci K., Ozdemir A., Hamaloglu E., Sokmener C. Perforation of acute calculus Meckel's diverticulitis: a rare cause of acute abdomen in elderly. *ActaChir. Belg.* 2000;226-227.
- [2]. Modi S, Pilai SK, DeClercq S. Perforated meckel's diverticulum in an adult due to fecolith: A rare case report and review of literature. *Int J Surg Case Rep.* 2015;15:143-5.
- [3]. Ding Y, Zhou Y, JiZh, Zhang J, Wang Q. Laparoscopic Management of Perforated Meckel's Diverticulum in Adults. *Int J Med Sci.* 2012;9(3):243-7.
- [4]. Hager M., Maier H., Eberwein M. Perforated Meckel's diverticulum presenting as a gastrointestinal stromal tumor: a case report. *J. Gastrointest. Surg.* 2005;9(6):809-811.
- [5]. Tas I, Culcu S, Duzkoylu Y, Eryilmaz S, Deniz MM, Yilmaz D. A rare cause of acute abdomen: Perforation of double Meckel's diverticulum. *Case Reports in Gastrointestinal Medicine.* 2015:648417.

- [6]. Farah RH et al. Spontaneous Perforation of Meckel's Diverticulum: A case report and review of literature. Pan Afr Med J. 2015;20:319.
- [7]. Quarrie R, Lindsey D, Bahner DP. Review of the Incidence and Management of Meckel's Diverticulum. Austin J Surg. 2014;1(3):1015.

Dr.Vineetkumar Mishra.“ Sponataneous perforation of Meckel's diverticulum: A Case report.”
IOSR Journal of Dental and Medical Sciences (IOSR-JDMS), vol. 18, no. 1, 2019, pp 47-49.