

## Weak Convergence Theorem of Khan Iterative Scheme for Nonself I-Nonexpansive Mapping

Poonam Lata Sagar<sup>1</sup>, S. K. Malhotra<sup>2</sup>

<sup>1</sup>Asst. Prof. Dept. of Applied Mathematics & Comp. Science, SATI (Deg), Vidisha (M.P.) India

<sup>2</sup>Head of dept. of Mathematics, Govt. S. G. S. P.G. College Ganj Basoda, Distt Vidisha (M.P.) India

**Abstract:** In this paper, we prove the weak convergence of a modified Khan iteration for nonself I-nonexpansive mapping in a Banach space which satisfies Opial's condition. Our result extends and improves these announced by S. Chornphrom and S. Phonin [Weak Converges Theorem of Noor iterative Scheme for Nonself I-Nonexpansive mapping, Thai Journal of Mathematics Volume 7(2009) no.2:311-317].

**Keywords:** Khan iterative scheme; weak convergence; nonself nonexpansive mapping; fixed point; Banach space.

**2000 Mathematics Subject Classification:** 47H09; 47H10 (2000 MSC)

### I. Introduction

Let  $E := (E, \|\cdot\|)$  be a real Banach space,  $K$  be a nonempty convex subset of  $E$ , and  $T$  be a self mapping of  $K$ . The Mann iteration [9] is defined as  $x_1 \in K$  and

$$x_{n+1} = (1 - \alpha_n)x_n + \alpha_n T x_n, \quad n \geq 1. \quad (1.1)$$

$$y_n = (1 - \beta_n)x_n + \beta_n T x_n$$

$$x_{n+1} = (1 - \alpha_n)x_n + \alpha_n T y_n, \quad n \geq 1. \quad (1.2)$$

The Noor iteration [8] is defined as  $x_1 \in K$  and  $z_n = (1 - \gamma_n)x_n + \gamma_n T x_n$

$$y_n = (1 - \beta_n)x_n + \beta_n T z_n$$

$$x_{n+1} = (1 - \alpha_n)x_n + \alpha_n T y_n, \quad n \geq 1, \quad (1.3)$$

The Khan iteration [9] is defined as  $x_1 \in K$  and

$$\begin{aligned} X_{n+1} &= (1 - \alpha_n)T^n x_n + \alpha_n S^n y_n \\ y_n &= (1 - \beta_n)x_n + \beta_n T^n x_n, \quad n \geq 1, \end{aligned} \quad (1.4)$$

Where  $\{\alpha_n\}, \{\beta_n\}, \{\gamma_n\} \subset (0, 1]$ .

In the above taking  $\beta_n = 0$  in (1.2) and taking  $\beta_n = 0, \gamma_n = 0$  in (1.3) we obtain iteration (1.1).

In 1975, Baillon [1] first introduced nonlinear ergodic theorem for general non-expansive mapping in a Hilbert space  $H$ : if  $K$  is a closed and convex subset of  $H$  and  $T$  has a fixed point, then for every  $x \in K, \{T^n x\}$  is a weakly almost convergent, as  $n \rightarrow \infty$ , to a fixed point of  $T$ . It was also

shown by Pazy [11] that if  $H$  is a real Hilbert space and  $(1/n) \sum_{i=0}^{n-1} T^i x$  converges weakly, as  $n \rightarrow \infty$ , to  $y \in K$ ,

$y \in F(T)$ .

In 1941, Tricomi introduced the concept of a quasi-nonexpansive mapping for real functions. Later Diaz and Metcalf [2] and Dotson [3] studied quasi-nonexpansive mappings in Banach spaces. Recently, this concept was given by Kirk [6] in metric spaces which we adapt to a normed space as the following:  $T$  is called a quasi-nonexpansive mapping provided for all  $x \in K$  and  $f \in F(T)$ .

$$\|Tx - f\| \leq \|x - f\| \quad (1.5)$$

Recall that a Banach space  $E$  is said to satisfy Opial's condition [10] if, for each sequence  $\{x_n\}$  in  $E$ , the condition  $x_n \rightarrow x$  implies that

$$\limsup_{n \rightarrow \infty} \|x_n - x\| < \limsup_{n \rightarrow \infty} \|x_n - y\| \tag{1.6}$$

for all  $y \in E$  with  $y \neq x$ . It is well known from [10] that all  $l_p$  spaces for  $1 < p < \infty$  have this property. However, the  $l_p$  spaces do not, unless  $p = 2$ . The following definitions and statements are needed for the proof of our theorem.

Let  $K$  be a closed convex bounded subset of uniformly convex Banach spaces  $E$  and  $T$  self-mapping of  $E$ . Then  $T$  is called nonexpansive on  $K$  if

$$\|Tx - Ty\| \leq \|x - y\| \tag{1.7}$$

for all  $x, y \in K$ . Let  $F(T) := \{x \in K : Tx = x\}$  be denote the set of fixed points of a mapping  $T$ .

Let  $K$  be a subset of a normed space  $E$  and  $T$  and  $I$  self-mappings of  $K$ . Then  $T$  is called  $I$ -nonexpansive on  $K$  if

$$\|Tx - Ty\| \leq \|Ix - Iy\| \tag{1.8}$$

for all  $x, y \in K$  [14].

A mapping  $T$  is called  $I$ -quasi-nonexpansive on

$$\|Tx - f\| \leq \|Ix - f\| \tag{1.9}$$

for all  $x, y \in K$  and  $f \in F(T) \cap F(I)$ .

A subset  $K$  of  $E$  is said to be a retract of  $E$  if there exists a continuous map  $P : E \rightarrow K$  such that  $Px = x$  for all  $x \in K$ . A map  $P : E \rightarrow E$  is said to be a retraction if  $P^2 = P$ . It follows that if a map  $P$  is a retraction, then  $Py = y$  for all  $y$  in the range of  $P$ . A set  $K$  is optimal if each point outside  $K$  can be moved to be closer to all points of  $K$ . Note that every nonexpansive retract is optimal. In strictly convex Banach spaces, optimal sets are closed and convex. However, every closed convex subset of a Hilbert space is optimal and also a nonexpansive retract.

**Remark 1.1.** From the above definitions it is easy to see that if  $F(T)$  is nonempty, a nonexpansive mapping must be quasi-nonexpansive, and linear quasi-nonexpansive mappings are nonexpansive. But it is easily seen that there exist nonlinear continuous quasi-nonexpansive mappings which are not nonexpansive. There are many results on fixed points on nonexpansive and quasi-nonexpansive mappings in Banach spaces and metric spaces. For example, Petryshyn and Williamson [12] studied the strong and weak convergence of the sequence of certain iterates to a fixed point of quasi-nonexpansive mapping. Their analysis was related to the convergence of Mann iterates studied by Dotson [3]. Subsequently, Ghosh and Debnath [4] considered the convergence of Ishikawa iterates of quasi-nonexpansive mappings in Banach spaces. Later Temir and Gul [15] proved the weakly convergence theorem for  $I$ -asymptotically quasi-nonexpansive mapping defined in Hilbert space. In [16], the convergence theorems of iterative schemes for nonexpansive mappings have been presented and generalized.

In [13], Rhoades and Temir considered  $T$  and  $I$  self-mappings of  $K$ , where  $T$  is  $I$ -nonexpansive mapping. They established the weak convergence of the sequence of Mann iterates to a common fixed point of  $T$  and  $I$ . More precisely, they proved the following theorems.

**Theorem (Rhoades and Temir [13]):** Let  $K$  be a closed convex bounded subset of uniformly convex Banach space  $E$ , which satisfies Opial's condition, and let  $T, I$  self-mappings of  $K$  with  $T$  be an  $I$ -nonexpansive mapping,  $I$  a nonexpansive on  $K$ . Then, for  $x_0 \in K$ , the sequence  $\{x_n\}$  of modified Noor iterates converges weakly to common fixed point of  $F(T) \cap F(I)$ .

In the above theorem,  $T$  remains self-mapping of a nonempty closed convex subset  $K$  of a uniformly convex Banach space. If, however, the domain  $K$  of  $T$  is a proper subset of  $E$  and  $T$  maps  $K$  into  $E$  then, the iteration formula (1.1) may fail to be well defined. One method that has been used to overcome this in the case of single operator  $T$  is to introduce a retraction  $P : E \rightarrow K$  in the recursion formula (1.1) as follows:  $x_1 \in K$ ,

$$x_{n+1} = (1 - \alpha_n)x_n + \alpha_n P T x_n, n \geq 1.$$

In [7], Kiziltunc and Ozdemir considered  $T$  and  $I$  are nonself mapping of  $K$  where  $T$  is an  $I$ -nonexpansive mapping. They established the weak convergence of the sequence of the modified Ishikawa iterative scheme to a common fixed point of  $T$  and  $I$ .

$$y_n = P((1 - \beta_n)x_n + \beta_n T x_n)$$

$$x_{n+1} = P((1 - \alpha_n)x_n + \alpha_n T y_n), \quad n \geq 1. \tag{1.10}$$

In this paper, we consider  $T$  and  $I$  are nonself mappings of  $K$ , where  $T$  is an  $I$ -nonexpansive mappings. We prove the weak convergence of the sequence of modified Noor iterative scheme to a common fixed point of  $F(T) \cap F(I)$ .

## II. Main Results

In this section, we prove the weak convergence theorem.

**Theorem 2.1.** Let  $K$  be a closed convex bounded subset of a uniformly convex Banach space  $E$  which satisfies Opial's condition, and let  $T, I$  nonself mappings of  $K$  with  $T$  be an  $I$ -nonexpansive mapping,  $I$  a nonexpansive on  $K$ . Then, for  $x_0 \in K$ , the sequence  $\{x_n\}$  of modified Khan iterates defined by  $x_1 \in K$ ,

$$\begin{aligned} Z_n &= (1 - \gamma_n)x_n + \gamma_n T^n x_n, \\ y_n &= (1 - \beta_n)x_n + \beta_n S_n Z_n, \\ x_{n+1} &= (1 - \alpha_n)x_n + \alpha_n P^n y_n, \end{aligned} \quad n \geq 1. \tag{2.1}$$

converges weakly to common fixed point of  $F(T) \cap F(I)$ .

**Proof.** If  $F(T) \cap F(I)$  is nonempty and a singleton, then the proof is complete. We will assume that  $F(T) \cap F(I)$  is nonempty and that  $F(T) \cap F(I)$  is not a singleton.

$$\begin{aligned} \|x_{n+1} - f\| &= \|(1 - \alpha_n)x_n + \alpha_n P^n y_n - f\| \\ &= \|(1 - \alpha_n)x_n + \alpha_n P^n y_n - (1 - \alpha_n + \alpha_n)f\| \\ &\leq (1 - \alpha_n)\|x_n - f\| + \alpha_n\|P^n y_n - f\| \\ &\leq (1 - \alpha_n)\|x_n - f\| + \alpha_n K_n \|y_n - f\| \end{aligned} \tag{2.2}$$

and

$$\begin{aligned} \|y_n - f\| &= \|(1 - \beta_n)x_n + \beta_n S^n z_n - f\| \\ &= \|(1 - \beta_n)x_n + \beta_n (S^n z_n - f)\| \\ &\leq (1 - \beta_n)\|x_n - f\| + \beta_n\|S^n z_n - f\| \\ &\leq (1 - \beta_n)\|x_n - f\| + \beta_n K_n \|z_n - f\| \end{aligned} \tag{2.3}$$

and also, we get

$$\begin{aligned} \|z_n - f\| &= \|(1 - \gamma_n)x_n + \gamma_n T^n x_n - f\| \\ &= \|(1 - \gamma_n)x_n + \gamma_n (T^n x_n - f)\| \\ &\leq (1 - \gamma_n)\|x_n - f\| + \gamma_n\|T^n x_n - f\| \\ &\leq (1 - \gamma_n)\|x_n - f\| + \gamma_n K_n \|x_n - f\| \end{aligned} \tag{2.4}$$

Substituting (2.4) in (2.3), we have

$$\|y_n - f\| \leq (1 - \beta_n)\|x_n - f\| + K_n \beta_n (1 - \gamma_n + K_n \gamma_n)\|x_n - f\| \tag{2.5}$$

Substituting (2.5) in (2.2), we have

$$\begin{aligned} \|x_{n+1} - f\| &\leq (1 - \alpha_n)\|x_n - f\| + \alpha_n K_n (1 - \beta_n + K_n \beta_n - K_n \beta_n \gamma_n + K_n^2 \beta_n \gamma_n)\|x_n - f\| \\ &\leq (1 - \alpha_n + K_n \alpha_n - K_n \alpha_n \beta_n + K_n^2 \alpha_n \beta_n - K_n^2 \alpha_n \beta_n \gamma_n + K_n^3 \alpha_n \beta_n \gamma_n)\|x_n - f\| \\ &\leq [1 - \alpha_n (K_n - 1) + \alpha_n \beta_n K_n (K_n - 1) + \alpha_n \beta_n \gamma_n K_n^2 (K_n - 1)]\|x_n - f\| \end{aligned}$$

Thus  $\alpha_n \neq 0, \beta_n \neq 0$  and  $\gamma \neq 0$ . Since  $\{K_n\}$  is a nonincreasing bounded sequence and hence  $K_n < 1$  implies that  $\sum_{n=1}^{\infty} (K_n - 1) < \infty$ . Then  $\lim_{n \rightarrow \infty} \|x_n - f\|$  exists.

Now we show that  $\{x_n\}$  converges weakly to a common fixed point of  $T$  and  $I$ . The sequence  $\{x_n\}$  contains a subsequence which converges weakly to a point in  $K$ . Let  $\{x_{nk}\}$  and  $\{x_{mk}\}$  be two subsequences of  $\{x_n\}$  which converge weak to  $f$  and  $q$ , respectively. We will show that  $f = q$ . Suppose that  $E$  satisfies Opial's condition and that  $f \neq q$  is in weak limit set of the sequence  $\{x_n\}$ . Then  $\{x_{nk}\} \rightarrow f$  and  $\{x_{mk}\} \rightarrow q$ , respectively. Since  $\lim_{n \rightarrow \infty} \|x_n - f\|$  exists for any  $f \in F(T) \cap F(I)$ , by opial's condition, we conclude that

$$\begin{aligned} \lim_{n \rightarrow \infty} \|x_n - f\| &= \lim_{k \rightarrow \infty} \|x_{nk} - f\| < \lim_{k \rightarrow \infty} \|x_{nk} - q\| \\ &= \lim_{n \rightarrow \infty} \|x_n - q\| = \lim_{j \rightarrow \infty} \|x_{mj} - q\| \\ &< \lim_{j \rightarrow \infty} \|x_{mj} - f\| = \lim_{n \rightarrow \infty} \|x_n - f\| \end{aligned}$$

This is a contradiction. Thus  $\{x_n\}$  converges weakly to an element of  $F(T) \cap F(I)$ . This completes the proof.

**Corollary 2.2.**(Kumam et al.[8, Theorem 2.1]) Let  $K$  be a closed convex bounded subset of a uniformly convex Banach space  $X$ , which satisfies Opial's condition, and let  $T, I$  self-mappings of  $K$  with  $T$  be an  $I$ -quasi-nonexpansive mapping,  $I$  a nonexpansive on  $K$ . Then, for  $x_0 \in K$ , the sequence  $\{x_n\}$  of three-step Noor iterative scheme defined by (1.3) converges weakly to common fixed point of  $F(T) \cap F(I)$ .

**Corollary 2.3.** (Kiziltunc and Ozdemir [7, Theorem 2.1]) Let  $K$  be a closed convex bounded subset of a uniformly convex Banach space  $E$ , which satisfies Opial's condition, and let  $T, I$  nonself mappings of  $K$  with  $T$  be an  $I$ -nonexpansive mapping,  $I$  a nonexpansive on  $K$ . Then, for  $x_1 \in K$ , the sequence  $\{x_n\}$  of modified Ishikawa iterates defined by (1.9) converges weakly to common fixed point of  $F(T) \cap F(I)$ .

**Theorem 24.** Let  $K$  be a closed convex bounded subset of a uniformly convex Banach space  $E$ , which satisfies Opial's condition, and let  $T, I$  nonself mappings of  $K$  with  $T$  be an  $I$ -nonexpansive mapping,  $I$  a nonexpansive on  $K$ . Then, for  $x_1 \in K$ , the sequence  $\{x_n\}$  of Mann converges weakly to common fixed point of  $F(T) \cap F(I)$ .

**Proof.** Putting  $\gamma_n = 0$  and  $\beta_n = 0$  in Theorem 2.1, we obtain the desired result.

### References

- [1] J. B. Baillon, Un theorem de type ergodique pour les contractions non lineaires dans un espace de Hilbert, Comptes Rendus de l'Academie des Sciences de Paris, Serie A 280(1975), no. 22, 1511-1514.
- [2] J. B. Diaz and F. T. Metcalf, On the set of sub sequential limit points of successive approximations, Transaction of the American Mathematical Society 135(1969), 459-485.
- [3] W. G. Dotson Jr., On the Mann iterative process, Transactions of the American Mathematical Society 149(1970), no. 1, 65-73.
- [4] M.K.Ghosh and L.Debnath, Convergence of Ishikawa iterates of quasi -nonexpansive mappings, Journal of Mathematical Analysis and Applications 207(1997), no. 1, 96-103.
- [5] S. Ishikawa, Fixed points by a new iteration method, Proc. Am. Math. Soc.44(1974) 147-150.
- [6] W. A. Kirk, Remarks on approximation and approximate fixed points in metric fixed point theory, Annales Universitatis Mariae Curie-Sklodowska. Sectio A 51(1997), no. 2, 167-178.
- [7] H. Kiziltunc and M. Ozdemir, On convergence theorem for nonself I- nonexpansive mapping in Banach spaces, Applied Mathematical Sciences 1(2007), no. 48, 2379-2383.
- [8] P. Kumam, W. Kumethong and N. Jewwaiworn, Weak convergence Theorems of three-step Noor iterative scheme for I-quasi-nonexpansive mappings in Banach spaces, Applied Mathematical Sciences 2(2008), no. 59, 2915-2920.
- [9] S.H.Khan, Y.J., M. Abbas, Convergence to common fixed points by a modified Iteration process, Journal of Appl. Math. andcomput. doi. 10. 1007/s 12190-010-0381-z.
- [10] Z. Opial, Weak convergence of the sequence of successive approximations for nonexpansive mappings, Bulletin of the American Mathematical Society 73(1967), 591-597.
- [11] A.Pazy, On the asymptotic behavior of iterates of nonexpansive mappings in Hilbert space, Israel Journal of Mathematics 26(1977), no. 2, 197-204.
- [12] W. V. Petryshyn and T. E. Williamson Jr., Strong and weak convergence of the sequence of successive approximations for quasi-nonexpansive mappings, Journal of Mathematical Analysis and Application 43(1973), 459-497.
- [13] B. E. Rhoades and S. Temir, Convergence theorems for I-nonexpansive mapping, to appear in International Journal of Mathematics and Mathematical Sciences.
- [14] N. Shahzad, Generalized I-nonexpansive maps and best approximations in Banach spaces, Demonstratio Mathematica 37(2004), no. 3, 597-600.
- [15] S. Temir and O. Gul, Convergence theorem for I-asymptotically quasi- nonexpansive mapping in Hilbert space, Journal of Mathematical Analysis and Applications 329(2007) 759-765.
- [16] H. Zhou, R. P. Agarwal, Y. J. Cho, and Y. S. Kim, Nonexpansive mappings and iterative method in uniformly convex Banach spaces, Georgian Mathematical Journal 9(2002), no. 3, 591-600.