

# Susceptible-Exposed-Infectious- Recovered-Undetectable- Susceptible (SEIRUS) Model Predictions of the Spread of COVID-19 Pandemic

Victor Alexander Okhuese<sup>1</sup>, Oduwole .H.K<sup>2</sup>, Jacob Emmanuel<sup>3</sup>

<sup>1,2</sup>Department of Mathematics, Nasarawa State University, Keffi – Nigeria

Department of Mathematics, University of Abuja, Abuja-Nigeria

## Abstract

The model equations which exhibits the disease-free equilibrium ( $E_0$ ) state for COVID-19 coronavirus does not exist and hence does not satisfy the criteria for a locally or globally asymptotic stability when the basic reproductive number  $R_0 = 1$  for and endemic situation. This implies that the COVID-19 coronavirus does not have a curative vaccine yet and precautionary measures are advised through quarantine and observatory procedures. The basic reproductive number was found to be  $R_0 < 1$  and hence shows that there is a chance of decline of secondary infections when the ratio between the incidence rate in the population and the total number of infected population quarantined with observatory procedure.

Furthermore, numerical simulations were carried to complement the analytical results in investigating the effect of the implementation of quarantine and observatory procedures has on the projection of the further spread of the virus globally. Result shows that the proportion of infected population in the absence of curative vaccination will continue to grow globally meanwhile the recovery rate will continue slowly which therefore means that the ratio of infection to recovery rate will determine the death rate that is recorded globally.

Therefore, the effort to evaluate the disease equilibrium shows that unless there is a dedicated effort from individual population, government, health organizations, policy makers and stakeholders, the world would hardly be reed of the COVID-19 coronavirus and further spread is eminent and the rate of infection will continue to increase despite the increased rate of recovery until a curative vaccine is developed.

**Keywords:** coronavirus pandemic globally, COVID-19 coronavirus, mathematical modeling of infection disease, SEIRUS-model, parameter identification, statistical methods.

Date of Submission: 05-07-2020

Date of Acceptance: 21-07-2020

## I. Introduction

A recent study by [1-2] focuses on the epidemic outbreak caused by COVID-19 coronavirus due to the global trend of the pandemic with its origin from mainland China. In his study, [1] used the popular SIR (Susceptible-Infectious-Removed) model to obtain optimal values for the model parameters with the use of statistical approach and hence predicated the number of infected, susceptible and removed persons versus time. This model approach by [1] has been a major breakthrough in modelling disease control as used by several authors ([2-3], among others). However, although there exist a global interest in knowing the rate of infection that will occur over time globally, it is of great interest to propose a mathematical model for the end in the spread and subsequent elimination of the virus. Hence, in this study we adopt solutions from [4] for a new deterministic endemic model (Susceptible – Exposed – Infectious – Removed – Undetectable – Susceptible: SEIRUS) originally developed for the control of the prevalence of HIV/AIDS in Africa.

The resulting equations are a system of coupled homogenous differential equations. Numerical experiments with relevant simulation showing how the variation of the reproductive number ( $R_0$ ) affect the number of infected individuals is carried out. Conscious effort through evaluating the new deterministic SEIRUS model is done to reduce the reproductive number ( $R_0$ ) to zero for a possible halt of the spread of the disease thereby leading to an endemic equilibrium to eradicate the disease in a later time in the future.

In summary, this study aims to develop and evaluate the new deterministic endemic SEIRUS compartmental model of the COVID-19 coronavirus dynamics which combines quarantine/observatory procedures and behavioral change/social distancing in the control and eradication of the disease in the most exposed sub-population.

The wide spread of the COVID-19 coronavirus and the lack or inefficiency of purposeful and result based intervention is a great call for other empirical and scientific interventions which seeks to review strategic models and recommendations of social and scientific research for disease control. Although previous studies

have been tailored towards the epidemiology and the disease-free equilibrium where the reproductive number of the infectious population is at its barest minimum, this study seeks to study the evaluative impact of the endemic equilibrium of a new endemic deterministic model while taking into consideration that possibility of the recovered population being Undetectable and fit to be moved to the Susceptible compartment which will therefore imply a zero secondary infection of the disease globally.

Also, this research work will contribute significantly to recent research findings by [1-2] as well as inform government, non-government organizations as well as policy maker's decision on sustainable policies to prevent the spread of the disease based on specific age groups of the active population.

**The Model Variables and Parameters**

The model variables and parameters for the investigation of the stability analysis of the equilibrium state for the new deterministic endemic model is given by;

<b>Variable</b>	<b>Description</b>
$S(t)$	Number of susceptible population at time $t$
$E(t)$	Number of exposed population at time $t$
$I(t)$	Number of infected population at time $t$
$R(t)$	Number of infected population quarantined and expecting recovery at time $t$
$U(t)$	Number of recovered adults satisfying undetectable criteria at time $t$

<b>Parameter</b>	<b>Description</b>
$\mu$	Natural death rate of the population
$\alpha_0$	Maximum death rate due to COVID-19 coronavirus. ( $\alpha \leq \alpha_0$ )
$\alpha$	Death rate of infected population due to COVID-19 coronavirus
$\varphi$	Disease induced death rate of infected population not quarantined
$\varpi$	Disease induced death rate of infected receiving quarantined
$T$	Maximum lifespan after infection ( $T \geq 14 \text{ days}$ )
$k$	Efficacy of Quarantine ( $0 \leq k \leq 1$ )
$\rho$	Rate of recovery
$\beta$	rate of transmission
$\sigma$	Proportion of infected population in quarantine per unit time (Treatment rate)
$\pi$	Proportion of population from susceptible to exposed/latent class
$\varepsilon$	Proportion of removed population still being observed and being moved to susceptible class.
$B(t)$	Incidence rate or force of infection in the population

**Model Assumptions**


The following assumptions would help in the derivation of the model:

1. There is no emigration from the total population and there is no immigration into the population. A negligible proportion of individuals move in and out of the population at a given time.
2. Maturation (or maturity) is interpreted as the period between infection to the period of symptoms observation (days 1 to 14)
3. The susceptible population are first exposed to a latent class where they can be infected or not.
4. Some infected individuals move to the removed class when they are quarantined for observatory procedures.
5. The recruitment from the  $S$ -class into the  $E$ -class is through contacts from population in the  $I$ -class to the  $S$ -class
6. The recruitment into the  $R$ -class from the  $I$ -class at a rate  $\sigma$ .
7. The recruitment into the  $U$ -class from the  $R$ -class depends on the effectiveness of the quarantine and observatory procedure at a rate  $\rho$ .
8. Death is implicit in the model and it occurs in all classes at constant rate  $\mu$ . However, there is an additional death rate in the  $I$  and  $R$  classes due to infection for both juvenile and adult sub-population denoted by  $\varphi$  and  $\varpi$  respectively.

**Model Description**

This study uses the deterministic endemic model where a susceptible class is a class that is yet to be infected, but is open to infection as interactions with members of the  $I$ -class continuous. An infected individual is one who has contracted the coronavirus and is at some stage of infection. A removed individual is one that is confirmed to have the virus with its expected symptoms and is under quarantine while following relevant observatory procedures. A member of the undetectable class is one that has been removed and does not secrete the virus anymore and has been satisfied by the WHO standard to be in the undetectable class.

The following diagram describes the dynamic of SEIRUS framework, and will be useful in the formulation of model equations.



**The Model Equations**

From the assumptions and the flow diagram above, the following model equations are derived.

$$\frac{dS(t)}{dt} = \pi + \mu S \tag{1}$$

$$\frac{dE(t)}{dt} = S - (\mu + B)E \tag{2}$$

$$\frac{dI(t)}{dt} = BE - (\sigma + \mu + \varphi)I \tag{3}$$

$$\frac{dR(t)}{dt} = \sigma I - (\rho + \mu + \omega)R \tag{4}$$

$$\frac{dU(t)}{dt} = \rho R - (\mu + \varepsilon)U \tag{5}$$

Such that

$$N(t) = S(t) + E(t) + I(t) + R(t) + U(t) \tag{6}$$

The incidence rate or force of infection at time  $t$  denoted by  $B(t)$  in the population is given as

$$B(t) = \frac{\beta I + \sigma R + \rho U}{N} \tag{7}$$

To simplify the model, we normalize the model by transforming the model equations into proportions. The derive model equations in proportion of infected juveniles and adults define prevalence of infection, which has biological meaning.

The model equations are transformed into proportions as follows;

$$\frac{dN(t)}{dt} = \frac{dS(t)}{dt} + \frac{dE(t)}{dt} + \frac{dI(t)}{dt} + \frac{dR(t)}{dt} + \frac{dU(t)}{dt} \tag{8}$$

$$\frac{dN(t)}{dt} = \pi + \mu S + S - \mu E - (\mu + \varphi)I - (\mu + \omega)R - (\mu + \varepsilon)U \tag{9}$$

Let

$$s = \frac{S}{N}, e = \frac{E}{N}, i = \frac{I}{N}, r = \frac{R}{N}, u = \frac{U}{N} \tag{10}$$

Then the normalized system is follows,

$$\frac{ds}{dt} = \frac{1}{N} \left[ \frac{dS(t)}{dt} - s \frac{dN(t)}{dt} \right]$$

Substituting (1) and (9) and using (10)

$$\frac{ds}{dt} = \frac{\pi(1-s)}{N} + \mu s - \mu s^2 - s s^2 + \mu s e + (\mu + \varphi) s i + (\mu + \omega) s r + (\mu + \varepsilon) s u \tag{11}$$

Similarly,

$$\frac{de}{dt} = \frac{1}{N} \left[ \frac{dE(t)}{dt} - e \frac{dN(t)}{dt} \right]$$

Substituting Equations (2) and (9) and using (10)

$$\frac{de}{dt} = s - (\mu + B)e - \frac{e\pi}{N} - \mu s e - e s + \mu e^2 + (\mu + \varphi) e i + (\mu + \omega) e r + (\mu + \varepsilon) e u \tag{12}$$

Next we have;

$$\frac{di}{dt} = \frac{1}{N} \left[ \frac{dI(t)}{dt} - i \frac{dN(t)}{dt} \right]$$

Substituting Equation (3) and (9) and using (10)

$$\frac{di}{dt} = B e - (\sigma + \mu + \varphi) i - \frac{i\pi}{N} - i \mu s - i s + \mu i e + (\mu + \varphi) i^2 + (\mu + \omega) i r + (\mu + \varepsilon) i u \tag{13}$$

And

$$\frac{dr}{dt} = \frac{1}{N} \left[ \frac{dR(t)}{dt} - r \frac{dN(t)}{dt} \right]$$

Substituting Equation (4) and (9) and using (10)

$$\frac{dr}{dt} = \sigma i - (\rho + \mu + \varpi)r - \frac{r\pi}{N} - \mu rs - rs + \mu re + (\mu + \varphi)ri + (\mu + \varpi)rr + (\mu + \varepsilon)ru \quad (14)$$

And finally;

$$\frac{du}{dt} = \frac{1}{N} \left[ \frac{dU(t)}{dt} - u \frac{dN(t)}{dt} \right]$$

Substituting Equation (5) and (9) and using (10)

$$\frac{du}{dt} = \rho r - (\mu + \varepsilon)u - \frac{u\pi}{N} - u\mu s - us + \mu ue + (\mu + \varphi)ui + (\mu + \varpi)ur + (\mu + \varepsilon)uu \quad (15)$$

However,

$$s + e + i + r + u = 1 \quad (16)$$

Equations (11) to (15) are the model equations in proportions, which define prevalence of infection.

**Existence and uniqueness of disease free equilibrium state ( $E_0$ ) of the SEIRUS model**

The disease-free equilibrium (DFE) state of the endemic SEIRUS model is obtained by setting the left hand sides of equations (11) – (15) to zero while setting the disease components  $e = i = r = u = 0$  leading to equations (17) – (18) below

$$0 = \frac{\pi(1-s)}{N} + \mu s - \mu s^2 - s s^2 \quad (17)$$

$$0 = s \quad (18)$$

And substituting (18) into (17) we have;

$$0 = \frac{\pi}{N}$$

Then taking (17) where  $s = 0$  or

$$0 = \mu - \mu s - s^2 \quad (19)$$

Simplifying further gives,

$$As^2 + Bs + C\mu = 0 \quad (20)$$

where  $A = 1, B = \mu$  and  $C = -\mu$

$$s^* = \frac{-\mu \pm \sqrt{\mu^2 + 4\mu}}{2} = \frac{-\mu \pm (\mu - 2)}{2}$$

Therefore, the solution for the simultaneous equations (19) is given by

$$(s_1^*, s_2^*) = \{(0,0), (-1, -\mu - 1)\} \quad (21)$$

Ignoring the native values of  $s_1^*$  and  $s_2^*$  and other stringent conditions, there exist a unique trivial and disease-free equilibrium states at  $(s_1^*, s_2^*)$  given by (0,0). The solution (21) satisfies equation (19) identically.

**Stability analysis of disease-free equilibrium state ( $E_0$ )**

In order to study the behavior of the systems (11) – (15) around the disease-free equilibrium state  $E_0 = [0,0,0,0,0]$ , we resort to the linearized stability approach such that;

$$f_s = \frac{\pi(1-s)}{N} + \mu s - \mu s^2 - s s^2 + \mu e + (\mu + \varphi)si + (\mu + \varpi)sr + (\mu + \varepsilon)su \quad (22)$$

$$f_e = s - (\mu + B)e - \frac{e\pi}{N} - \mu se - es + \mu e^2 + (\mu + \varphi)ei + (\mu + \varpi)er + (\mu + \varepsilon)eu \quad (23)$$

$$f_i = Be - (\sigma + \mu + \varphi)i - \frac{i\pi}{N} - i\mu s - is + \mu ie + (\mu + \varphi)i^2 + (\mu + \varpi)ir + (\mu + \varepsilon)iu \quad (24)$$

$$f_r = \sigma i - (\rho + \mu + \varpi)r - \frac{r\pi}{N} - \mu rs - rs + \mu re + (\mu + \varphi)ri + (\mu + \varpi)rr + (\mu + \varepsilon)ru \quad (25)$$

$$f_u = \rho r - (\mu + \varepsilon)u - \frac{u\pi}{N} - u\mu s - us + \mu ue + (\mu + \varphi)ui + (\mu + \varpi)ur + (\mu + \varepsilon)uu \quad (26)$$

Therefore;

$$\begin{aligned} \frac{\partial f_s}{\partial s} &= -\frac{\pi}{N} + \mu, \frac{\partial f_s}{\partial e} = 0, \frac{\partial f_s}{\partial i} = 0, \frac{\partial f_s}{\partial r} = 0, \frac{\partial f_s}{\partial u} = 0 \\ \frac{\partial f_e}{\partial s} &= 1, \frac{\partial f_e}{\partial e} = -(\mu + B) - \frac{\pi}{N}, \frac{\partial f_e}{\partial i} = 0, \frac{\partial f_e}{\partial r} = 0, \frac{\partial f_e}{\partial u} = 0 \\ \frac{\partial f_i}{\partial s} &= 0, \frac{\partial f_i}{\partial e} = B, \frac{\partial f_i}{\partial i} = -(\sigma + \mu + \varphi) - \frac{\pi}{N}, \frac{\partial f_i}{\partial r} = 0, \frac{\partial f_i}{\partial u} = 0 \\ \frac{\partial f_r}{\partial s} &= 0, \frac{\partial f_r}{\partial e} = 0, \frac{\partial f_r}{\partial i} = \sigma, \frac{\partial f_r}{\partial r} = -(\rho + \mu + \varpi) - \frac{\pi}{N}, \frac{\partial f_r}{\partial u} = 0 \end{aligned}$$

$$\frac{\partial f_u}{\partial s} = 0, \frac{\partial f_u}{\partial e} = 0, \frac{\partial f_u}{\partial i} = 0, \frac{\partial f_u}{\partial r} = \rho, \frac{\partial f_u}{\partial u} = -(\mu + \varepsilon) - \frac{\pi}{N}$$

And the Jacobian  $J_{E_0}$  is given by

$$J_{E_0} = \begin{bmatrix} -\frac{\pi}{N} + \mu & 0 & 0 & 0 & 0 \\ 1 & -(\mu + B) - \frac{\pi}{N} & 0 & 0 & 0 \\ 0 & B & -(\sigma + \mu + \varphi) - \frac{\pi}{N} & 0 & 0 \\ 0 & 0 & \sigma & -(\rho + \mu + \varpi) - \frac{\pi}{N} & 0 \\ 0 & 0 & 0 & \rho & -(\mu + \varepsilon) - \frac{\pi}{N} \end{bmatrix} \quad (27)$$

Hence, according to ([5], p255), the Determinant of the Jacobian Matrix  $J_{E_0}$  is given by the recursive definition for a  $5 \times 5$  matrix defined as;

$$Det(J_{E_0}) = a_{11}det(J_{E_011}) - a_{12}det(J_{E_012}) + a_{13}det(J_{E_013}) - \dots \pm a_{55}det(J_{E_055}) \quad (28)$$

And from (27)

$$Det(J_{E_0}) = 0 \quad (29)$$

Similarly from the Trace of the Jacobian matrix ( $J_{E_0}$ ) given in equation (27) we have that

$$Trace(J_{E_0}) = -\frac{5\pi}{N} - B - (\sigma + \mu + \varphi) - (\rho + \mu + \varpi) - (\mu + \varepsilon) > 0 \quad (30)$$

Hence, since  $Def(J_{E_0}) = 0$  and  $Trace(J_{E_0}) > 0$  which does not satisfy the prescribed threshold criteria based on [5], then the disease free equilibrium ( $E_0$ ) for COVID-19 coronavirus does not satisfy the criteria for a locally or globally asymptotic stability. This implies that as a pandemic as declared by [6] the COVID-19 coronavirus does not have a curative vaccine yet and precautionary measures are advised through quarantine and observatory procedures.

### Computation of the basic reproductive number ( $R_0$ ) of the model

The Basic Reproductive number ( $R_0$ ) is define as the number of secondary infections that one infectious individual would create over the duration of the infectious period, provided that everyone else is susceptible.  $R_0 = 1$  is a threshold below which the generation of secondary cases is insufficient to maintain the infection in human community. If  $R_0 < 1$ , the number of infected individuals will decrease from generation to next and the disease dies out and if  $R_0 > 1$  the number of infected individuals will increase from generation to the next and the disease will persist.

To compute the basic reproductive number ( $R_0$ ) of the model (22) – (26), we employ the next generation method as applied by [7,8,3].

$$\mathcal{F}_i = \begin{pmatrix} Be + \mu i \\ Be \end{pmatrix} \quad (31)$$

and

$$\mathcal{V}_i = \begin{pmatrix} \frac{\pi(1-s)}{N} + \mu s - \mu s^2 - s s^2 + \mu s e + (\mu + \varphi) s i + (\mu + \varpi) s r + (\mu + \varepsilon) s u \\ s - (\mu + B) e - \frac{e\pi}{N} - \mu s e - e s + \mu e^2 + (\mu + \varphi) e i + (\mu + \varpi) e r + (\mu + \varepsilon) e u \\ Be - (\sigma + \mu + \varphi) i - \frac{i\pi}{N} - i \mu s - i s + \mu i e + (\mu + \varphi) i^2 + (\mu + \varpi) i r + (\mu + \varepsilon) i u \\ f_r = \sigma i - (\rho + \mu + \varpi) r - \frac{r\pi}{N} - \mu r s - r s + \mu r e + (\mu + \varphi) r i + (\mu + \varpi) r r + (\mu + \varepsilon) r u \\ \rho r - (\mu + \varepsilon) u - \frac{u\pi}{N} - u \mu s - u s + \mu u e + (\mu + \varphi) u i + (\mu + \varpi) u r + (\mu + \varepsilon) u u \end{pmatrix} \quad (32)$$

where  $\mathcal{F}_i$  and  $\mathcal{V}_i$  are the rate of appearances of new infections in compartment  $i$  and the transfer of individuals into and out of compartment  $i$  by all means respectively. Using the linearization method, the associated matrices at disease-free equilibrium ( $E_0$ ) and after taking partial derivatives as defined by

$$D\mathcal{F}_i(E_0) = \begin{pmatrix} F & 0 \\ 0 & 0 \end{pmatrix} \text{ and } D\mathcal{V}_i(E_0) = \begin{pmatrix} V & 0 \\ J_1 & J_2 \end{pmatrix}$$

where  $F$  is nonnegative and  $V$  is a non-singular matrix, in which both are the  $m \times m$  matrices defined by

$$F = \left[ \frac{\partial \mathcal{F}_i}{\partial x_i}(E_0) \right]$$

and

$$V = \left[ \frac{\partial \mathcal{V}_i}{\partial x_i} (E_0) \right]$$

with  $1 \leq i, j \leq m$  and  $m$  is the number of infected classes.

In particular  $m = 2$ , we have

$$F = \begin{pmatrix} B & \mu \\ 0 & B \end{pmatrix}$$

and

$$V = \begin{pmatrix} -(\sigma + \varphi) & 0 \\ 0 & -(\sigma + \varphi) \end{pmatrix}$$

If the inverse of  $V$  is given as

$$V^{-1} = V = \begin{pmatrix} -\frac{1}{\sigma + \varphi} & 0 \\ 0 & -\frac{1}{\sigma + \varphi} \end{pmatrix}$$

Then the next matrix denoted by  $FV^{-1}$  is given as

$$FV^{-1} = \begin{pmatrix} -\frac{B}{\sigma + \varphi} & 0 \\ 0 & -\frac{B}{\sigma + \varphi} \end{pmatrix}$$

We find the eigenvalues of  $FV^{-1}$  by setting the determinant  $|FV^{-1} - \gamma I| = 0$

$$|FV^{-1} - \gamma I| = \begin{vmatrix} -\frac{B}{\sigma + \varphi} - \gamma & 0 \\ 0 & -\frac{B}{\sigma + \varphi} - \gamma \end{vmatrix} = 0$$

with characteristics polynomial

$$\rho(\gamma) = \gamma^2 + \left(\frac{2B}{\sigma + \varphi}\right)\gamma + \left(\frac{B}{\sigma + \varphi}\right)^2$$

and characteristics equation given as

$$\gamma^2 + \left(\frac{2B}{\sigma + \varphi}\right)\gamma + \left(\frac{B}{\sigma + \varphi}\right)^2 = 0$$

Solving the characteristics equation for the eigenvalues  $\gamma_{1,2}$ , where  $R_0$  is the maximum of the two eigenvalues  $\gamma_{1,2}$ . Hence the Basic Reproductive number is the dominant eigenvalues of  $FV^{-1}$ . Thus we have that

$$R_0 = -\left(\frac{B}{\sigma + \varphi}\right) < 1 \tag{33}$$

The Basic Reproductive number ( $R_0 < 1$ ) by Equation (33) shows that there is a chance of decline of secondary infections when the ratio between the incidence rate in the population and the total number of infected population quarantined with observatory procedure. Hence although there currently exist no clinical vaccine for the cure of COVID-19 coronavirus, with Equation (33) there is a high chance of decline in infection when all precautionary measures are observed globally.


## II. Results

### Description and Validation of Baseline Parameters for Global cases of COVID-19 coronavirus


According to the [9], the total cases of COVID-19 coronavirus worldwide stands: 142,320 and the current total deaths is 5,388 from about 129 countries.



**Figure 1:** A World Map showing the number of cases per each country with a COVID-19 coronavirus case. *Source: [9].*



**Figure 2:** A Cumulative Case chart showing the number of cases with a COVID-19 coronavirus case. *(Source: [9]).*



**Figure 3:** A Case by date of report chart showing the number of cases with a COVID-19 coronavirus case. *Source: [8].*

### III. Numerical Experiments of the Model

The age-structured deterministic model (11) – (15) was solved numerical using Runge-Kutta-Fehlberg 4-5th order method and implemented using Maple Software. The model equations were first transformed into proportions, thus reducing the model equations to ten differential equations. The parameters used in the


implementation of the model are shown in Table 1 below. Parameters were chosen in consonance with the threshold values obtained in the stability analysis of the disease free equilibrium state of the model.

**Table 1:** Estimated values of the parameters used in the Numerical experiments

Parameters	Values	Data Source	Parameters	Values	Data Source
$N(0)$	7.57bn	[10]	$\varphi$	0.000005*	Assumed
$N(1)$	149,292	[9]	$\omega$	0.0000007	[12]
$s(0)$	1.0000	Estimation	$T$	14 days	[9]
$e(0)$	1.0000	Estimation	$k$	0.5*	Assumed
$i(0)$	0.00002	[9]	$\rho$	0.000095	[12]
$r(0)$	0.000095	[12]	$\beta$	0.00002	[9]
$u(0)$	0.000095	[12]	$\sigma$	0.28404**	Estimated
$\mu$	0.000001	[10]	$\pi$	0.00567*	Assumed
$\alpha_0$	0.000011	[11]	$\varepsilon$	0.000095	[12]
$\alpha$			$B(t)$	0.0027**	Assumed

**Assumed\***: Hypothetical data use for research purpose **Assumed\*\***: Based on [1].

Hence from Equation (33) the Reproductive Number  $R_0 = -0.009505 < 1$  means there is a 99% chances of secondary infection when an infected population interact by contact with the susceptible population.



**Figure 4:** A chart showing the rate of recovery and infection from 1 to 14 days of the global COVID-19 coronavirus pandemic.

#### IV. Discussion

Figure 4 clearly shows that over a short period of time the rate of infection of the COVID-19 coronavirus will increase globally before normalizing at a point where future occurrence will be halted due to active quarantine and observatory procedures as prescribed by the [6] with the hope that a vaccine will be made ready in a shortest future possible.

Similarly, the chart also reveals that the rate of recovery will continue to increase despite the negligible increase in death rate from the COVID-19 coronavirus globally.

## V. Conclusions

In this study, the  $Def(J_{E_0}) = 0$  and  $Trace(J_{E_0}) > 0$  does not satisfy the prescribed threshold criteria based on Gerald (2012), then the disease free equilibrium ( $E_0$ ) for COVID-19 coronavirus does not satisfy the criteria for a locally or globally asymptotic stability. This implies that as a pandemic as declared by WHO (2020) the COVID-19 coronavirus does not have a curative vaccine yet and precautionary measures are advised through quarantine and observatory procedures. Also, the Basic Reproductive number ( $R_0 < 1$ ) by Equation (33) shows that there is a chance of decline of secondary infections when the ratio between the incidence rate in the population and the total number of infected population quarantined with observatory procedure.

The effort to evaluate the disease equilibrium shows that unless there is a dedicated effort from government, decision makers and stakeholders, the world would hardly be reed of the COVID-19 coronavirus and further spread is eminent. Therefore, as corroborated by the studies of [1-2], the rate of infection will continue to increase despite the increased rate of recovery because of the absence of vaccine at the moment.

Meanwhile, with observatory procedures and quarantine the rate of recovery will not only be enhanced, the chances of total recovery where the virus nor it symptoms will be undetectable is achievable as the rate of detection becomes negligibly small in the bodies of the recovered population.

### Declaration Statements:

Availability of data and material: The data used for this study was made available from; John Hopkins University (2020) for the total number of infected population and the recovered class of the entire population. Also, the data for the global population used to extract the susceptible class was made available from World Population Review (2019).

### Reference

- [1]. Nesteruk, I. (2020). Statistics based predictions of coronavirus 2019-nCoV spreading in mainland China. *MedRxiv*.
- [2]. Ming, W. K., Huang, J., & Zhang, C. J. (2020). Breaking down of healthcare system: Mathematical modelling for controlling the novel coronavirus (2019-nCoV) outbreak in Wuhan, China. *bioRxiv*.
- [3]. Oduwole, H. K. and Kimbir, A. R. (2018). "Modelling vertical transmission and the effect of Antiretroviral Therapy (ART) on the dynamics of HIV/AIDS in an age-structured population in Nigeria. *Journal of Natural and Applied Science-NasaraScientifique*, Vol. 7 No. 1 pp. 51 – 78.
- [4]. Victor, A. O. and Oduwole, H. K. (2020).Evaluating the deterministic SEIRUS mode for disease control in an age-structured population. *Global Scientific Journal*. Vol. 8. Issue 3.
- [5]. Gerald T. (2012). Ordinary Differential Equations and Dynamical Systems. *American Mathematical Society*. URL: <http://www.mat.univie.ac.at/~gerald/>
- [6]. WHO (March 11, 2020) WHO characterizes COVID-19 as a pandemic. Accessed by March 14, 2020 on: <https://www.who.int/emergencies/diseases/novel-coronavirus-2019/events-as-they-happen>
- [7]. Diekmann, O, Heesterbeek, J. A. P and Roberts, M. G. (2009); The construction of next-generation matrices for compartmental epidemic models, *J. R. Soc. Interface*, 2009 (doi:10.1098/rsif.2009.0386)
- [8]. Van den Driessche, P. and Watmough, J. (2002); Reproduction numbers and sub- threshold endemic equilibria for compartmental models of disease transmission. *Math. Biosci.* 180, 29–48, 2002 doi:10.1016/S0025-5564(02)00108-6
- [9]. WHO (March 13, 2020).Novel Coronavirus (COVID-19) Situation. Accessed by March 14, 2020 on: <https://experience.arcgis.com/experience/685d0ace521648f8a5beeee1b9125cd>
- [10]. World Population Review (WPR) (2020) - United Nations population estimates and projections. Total population: Estimated to be consistent with the 1963, 1991 and 2006 censuses, adjusted for underenumeration, with the age and sex structure from the 2011 MICS4 survey, and with estimates of the subsequent the trends in fertility, mortality and international migration.
- [11]. Batista, M. (2020). Estimation of the final size of the coronavirus epidemic by the SIR model. *ResearchGate*.
- [12]. John Hopkins University (JHU) (2020). Coronavirus COVID-19 Global Cases by the Center for Systems Science and Engineering (CSSE) at Johns Hopkins. Accessed by March 14, 2020 on: <https://www.arcgis.com/apps/opsdashboard/index.html#/bda7594740fd40299423467b48e9ecf6>

Victor Alexander Okhuese, et. al. "Susceptible-Exposed-Infectious- Recovered-Undetectable-Susceptible (SEIRUS) Model Predictions of the Spread of COVID-19 Pandemic." *IOSR Journal of Mathematics (IOSR-JM)*, 16(4), (2020): pp. 52-60.