

## Herbicidal potential of agrochemicals of *Sclerotium rolfsii* against *Parthenium hysterophorus*

<sup>1</sup>Dr Ramanuj Patel\* Deepika Patel\* and A.K. Pandey<sup>3</sup>

<sup>1</sup>Department of Biological Sciences, R.D. University, Jabalpur- 482001

<sup>2</sup>Department of Zoology and Biotechnology research lab Government Sciences College Jabalpur

<sup>3</sup>Chairman Private University Regulatory Commission Board Bhopal M.P

---

**Abstract:** Herbicidal potential of CFCF (Cell free culture filtrate) of *Sclerotium rolfsii* against *Parthenium hysterophorus* was determined by shoot cut, seedling and detached leaf bioassays. Maximum phytotoxicity was obtained from 14 day's old fermented broth with Sucrose + Tween 20 @0.5% as formulating agent. Significant reduction in chlorophyll and protein contents of host was also noticed.

**Key Words:** Herbicidal potential, *Sclerotium rolfsii*, *Parthenium hysterophorus*, Biorational Phytotoxicity, Formulation

---

### I. Introduction

*Parthenium hysterophorus* is an obnoxious, pernicious and deadly weed of compositae family native of North America, probably introduced in India along with grains under the PL 480 scheme. It poses serious threat to crops, livestock and human beings. Its infestation can be observed along railway lines, forest land, grassland and even valuable lands too. It is responsible for the substantial losses to the crops. It reduces agricultural yield by 40% and forage production by 90% (Knox et al., 2006). It has now gained the status of a major weed in forest areas of some states of India. It is gradually invading forest and other valuable lands. Conventional methods of its management rely mainly on the use of chemical herbicides. Public concern over the safety due to indiscriminate use of synthetic herbicides has generated significant pressures on weed scientists to search an alternative of these chemicals. Exploitation of microorganisms and especially their biorationals (natural products) as herbicides have generated significant interest world wide. (Pandey et al., 2002-2005). Survey conducted at various habitats of central India for weed pathogens yielded an isolate of *Sclerotium rolfsii* FGCC#02 which incites severe collar rot disease in *Parthenium* (Pandey et al., 1996a). Mycoherbicidal potential of the pathogen is known to influence by environmental factors.

To overcome these constraints, secondary metabolites especially oxalic acid synthesized by the pathogen have also tried (Pandey et al., 2003). Oxalic acid produced by the pathogen showed high herbicidal potential against *Parthenium*. Normal application of oxalic acid did not produced noticeable results and need a suitable formulating agent for its effective herbicidal potential. Therefore, the present investigation deals with the formulation and in vitro evaluation of herbicidal potential of the pathogen against *Parthenium* by Seedling and Detached leaf bioassay.

### II. Material And Methods

#### Recovery of Strains

*S.rolfsii* FGCC#02 was obtained from Fungal Germplasm Collection Centre, R.D.University, Jabalpur, previously isolated from the diseased part of the target weed and maintained on Potato Dextrose Agar (PDA) medium as per Agrawal and Hasija (1986).

#### Preparation and Extraction of CFCF

1000 ml Erlenmeyer flasks containing 500 ml of sterilized Richard's broth (KNO<sub>3</sub>-10g; MgSO<sub>4</sub>-2.5g; KH<sub>2</sub>PO<sub>4</sub>-2.5g; Sucrose-30g and Distilled Water-1000ml) were seeded with 5mm disc obtained from 7 days old cultures grown on PDA medium at 30<sup>o</sup> C. Inoculated flasks were incubated at 30 ± 2<sup>o</sup>C for 7 and 14 days. Fermented broth was filtered through Whatman's filter paper No.1 and the filtrate was centrifuged at 4000 rpm for 10 min in a centrifuge (Remi). Supernatant was discarded and the crude filtrate was again passed through 0.25µm Sartorius filter in vacuo condition (Abbas et al., 1992). Oxalic acid production was also determined by the method suggested by Bateman and Beer (1965). Herbicidal potential of the agrochemicals of 7 and 14 days old was evaluated by detached leaf and seedling bioassay (whole plant bioassay).

#### Detached leaf bioassay

Leaves from the host target weed were surface sterilized with 2% NaOCl and were kept on a sterilized moisture chamber prepared by using cotton and filter paper in a Petridish. (Thapar et al., 2002). Various

dilutions of toxic metabolites (5ml) were used of 7 and 14 days old fermented broth viz., 25%, 50%, 75% and 100% and the effects were observed after 24 hrs. All the experiments were performed in triplicates.

**Seedling bioassay**

Seedlings were raised in pots (10cm) containing soil, sand and peat in 1:1:1 ratio. Different concentration of S.rolfsii toxin (7 and 14 days old) was sprayed on host seedlings @ 30ml per plant and maintained in a green house (Temperature 30<sup>0</sup>C& Relative humidity 85-90%). Each treatment was replicated thrice with two control and observations were recorded after 3 days based on a score chart and percent disease index was calculated by using the following formula (Praveena 2003).

$$\text{Percent Disease Index} = \frac{\text{Sum of score each leaf}}{\text{No. of leaves scored} \times \text{Maximum score}} \times 100$$

**Effect of CFCF on chlorophyll and protein contents**

1gm of fresh leaves were homogenized with excess of ethanol in a mortar with pestle and centrifuged at 8000 rpm for 2min, with 80% ethanol. Supernatant was taken in other flask and diluted with ethanol. Absorbance of the extract was measured by UV- Vis Systronics spectrophotometer at 645 and 663nm for the determination of Chl a, Chl b and total Chl. The protein content was determined as per Lowry et al (1951).

**Formulation**

To test the compatibility of the toxin (14 days old) synthesized by the pathogen a total of 15 formulating agents namely Sorbitol, T-20+water, T-20+Sucrose, H<sub>2</sub>O + Gelatin, Toxin +Triton X 100, Toxin+ water, T-80+ sucrose+H<sub>2</sub>O, Cween 80, Toxin, Toxin + Acrylamide, Toxin+ Glycerol, Toxin+ coconut oil, Toxin+Soyabean, Toxin+Mustard oil, Toxin+Tween 80 were tried. All the formulating agents were added at rate of 0.5% to the toxin and its herbicidal potential was determined by seedling assay and observations were made after 48hrs up to 10<sup>th</sup> day. (Daigle and Conick 2002).

**III. Figures And Tables**

**Table: 1 Effect of different concentration of toxin on host seedlings**

Incubation period		
%Disease index		
Concentration	7 days	14 days
25 %	25±0.01	36±0.04
50 %	59±0.04	80 ± 0.02
75 %	50±0.25	75 ±0.25
100 %	48±0.01	73±0.15
Control	No effect	No effect

Amount of agrochemical sprayed: 30ml/plant; RH: 85%; Values given in the table are mean ± SEM

**Table: 2 Effect of different concentration of toxin on detached leaves of Host**

Incubation period		
%Disease index		
Concentration	7 days	14 days
25 %	33±0.01	55±0.01
50 %	66±0.05	95 ± 0.25
75 %	52±0.02	78 ±0.03
100 %	49±0.01	70±0.01
Control	No effect	No effect

Values given in the table are mean ± SEM; Amount of agrochemical used: 5ml/plant; RH: 85

**Table: 3 Effect of different concentration of toxin on biological contents of host seedlings**

Concentration	Biological Activity of CFCF							
	7 Days				14 Days			
	Chl.a	Chl.b	Total Chl.a	Protein	Chl.a	Chl.b	Total Chl.a	Protein
25 %	10.4±0.25	12.1±0.06	32.6±0.32	36.1±0.04	50.3±0.02	40.8±0.02	45.9±0.2	35.5±0.04
50 %	72.1±0.07	58.4±0.07	65.4±0.28	56.9±0.07	97.8±0.01	74.7±0.02	72.5±0.07	76.1±0.01
75 %	68.8±0.04	56.1±0.65	60.3±0.62	36.9±0.04	82.2±0.25	68.5±0.01	65.3±0.2	54.1±0.01
100 %	69.6±0.05	53.4±0.025	59.9±0.73	27.8±0.07	78.1±0.01	65.2± .03	63.4±0.3	42.2±0.02
Control	No effect	No effect	No effect	No effect	No effect	No effect	No effect	No effect

Values given in the table are mean ± SEM; Amount of agrochemical sprayed: 30ml/plant; RH: 85%

**Table: 4 Effect of different formulations on host seedlings under green house conditions**

SNo	Formulating agent	% Disease intensity				
		2 <sup>nd</sup> day	4 <sup>th</sup> day	6 <sup>th</sup> day	8 <sup>th</sup> day	10 <sup>th</sup> day
1	Sorbitol	1	4	4	4	4
2	T-20+water	2	5	5	5	5
3	T-20+Sucrose	5	6	6	6	6
4	H <sub>2</sub> O + Gelatin	5	5	5	5	5
5	Toxin +Triton X 100	2	5	5	5	5
6	Ioxin+ water	5	5	6	6	6
7	T-80+ sucrose+H <sub>2</sub> O	4	3	4	5	5
8	Cween 80	2	4	4	4	4
9	Toxin	6	6	6	6	6
10	Toxin + Acrylamide	2	4	4	4	4
11	Toxin+ Glycerol	1	1	1	1	1
12	Toxin+coconut oil	1	1	1	1	1
13	Toxin+Soyabean	3	3	3	3	3
14	Toxin+Mustard oil	3	3	3	4	4
15	Toxin+Tween 80	6	7	8	9	9
16	Control (Richard'sbroth)	No Effect				
	SEM	±0.33				
	CD <sub>5%</sub>	0.946				

Disease rating index (Horsfall and Barret 1945)

1 =99%,2=95%,3=91%,4=82%,5=62%,6=38%,7=18%,8=9%,9=5%,10=1%)

Amount of agrochemical sprayed: 30ml/plant; RH: 85%; Incubation period: 14 days

#### IV. Results And Conclusion

##### Seedling bioassay

It is clearly evident from the data recorded in **Table 1** that CFCF obtained from different incubations has varied effect on host seedlings. Maximum average leaf area damage (ALAD) of 85% on the seventh day and 65% of leaf damage by fourth day was observed when treated with CFCF obtained from 14 day's old fermented broth at 50% concentration followed by 75 and 100%. CFCF obtained from 14 days old fermented broth showed the maximum toxicity due to the maximum production of oxalic acid. CFCF obtained from 7 days old fermented broth also showed considerable toxicity at higher concentration. Similar observations have also reported by many other workers. (Pandey et al., 2002; Winder and Watson 1994; Saxena and Pandey 2001; Saxena et al., 2001) In contrast to this, several workers recorded maximum toxin production after 7 days of incubation.(Pandey et al., 2003) While the CFCF obtained from 14 d old fermented broth showed the maximum mortality to the host seedlings, 7 days old broth didn't cause significant damage to the host seedlings.. Similar findings have also been made by other workers. (Abbas et al., 1995; Shukla and Pandey 2006). Each experiment was performed in triplicates. The means of experiment was analyzed by one way ANOVA which indicates significant difference between the mean leaf area damage at different concentration with a P value of 5%.

##### Detached leaf assay

Noticeable symptoms were also observed when detached leaves were treated with CFCF of 7 and 14 days old fermented broth at various concentrations viz., 25, 50, 75 and 100%. Leaves were completely killed at 100% concentration and more than 80% of damage was recorded at 50% concentration (**Table 2**). Results obtained by detached leaf assay were quite prompting to use this pathogen as herbicidal agent against the weed. Similar damage ratings were also recorded by other workers (Joseph et al., 2002). One way ANOVA which indicates significant difference between the mean leaf area damage at different concentration of toxin with a P value of 5%.

##### Effect on chlorophyll and protein contents

Chlorophyll and protein contents were also significantly affected when treated with CFCF. The contents were gradually decreased with increased incubation. Maximum reduction was recorded in leaves treated with CFCF obtained from 14 days old fermented broth at 50% concentration followed by 75% and 100% (**Table 3**). 14 d old fermented broth showed the maximum biological activity as depicted by the significant reduction in the chlorophyll and protein content of the host leaves. While extract obtained from 7 days old broth failed to show any remarkable reduction in these contents at similar concentration. The effect was comparatively more on chlorophyll a and total chlorophyll while chlorophyll b and protein contents were less affected. Variation in toxicity in relation to incubation period may be due to different phase of growth of the fungus. Metabolites required for own growth are normally synthesized during initial phase whereas most of the toxicants

are formed during idiophase i.e. stationary phase of the fungus. (Abbas et al., 1995) also recorded 25 to 78% reduction in chlorophyll content in *Datura* sp. (jimson weed) tissues treated with fumonisin. Similarly significant biological activity of CFCF of many other microorganism including fungi have also recorded by several workers. (Sharma et al., 2004; Thapar et al., 2002; Abbas et al., 1995; Joseph et al., 2002; Saxena and Pandey 2001; Saxena et al., 2001 ; Kovics et al., 2005; Pandey and Pandey 2005).

### Formulation

Data recorded in **Table.4** clearly depicts that maximum phytotoxicity was shown by the toxin formulated with Sucrose + Tween 20 followed by Tween-80.Triton X was found to be highly inhibiting in its action. Rest of the agents produced average effect on the host shoots. Findings obtained in the above study clearly revealed the herbicidal potential of the pathogen against *Parthenium*. Maximum toxicity obtained at 50% concentration with sucrose +Tween 20 indicates the toxin compatibility with the formulating agent. However, others did not produce significant damage to the host seedling. Variations in different formulation may be due to the compatibility of the organism with various formulating agent. Similar findings have also been made by Singh (2002).

The above findings clearly indicate that the present isolate have significant potential to produce phytotoxic compounds with high herbicidal properties against *P. hysterophorus*. However, detailed investigation regarding characterization, standardization of large scale production of herbicidal compounds are to be carried out before its field application.

### Acknowledgement

We are grateful to Head Department of Biological Sciences, R.D.University, and Jabalpur for providing necessary laboratory facilities.

### References

- [1]. A.K. Pandey., J. Mishra, R.C. Rajak and S.K. Hasija, (1996a), Potential of indigenous strains *Sclerotium rolfsii* Sacc. For the management of *Parthenium hysterophorus* L. a serious threat to biodiversity in India. In: Herbal Medicines, Biodiversity and Conservation Strategies (R.C. Rajak & M.K. Rai eds), International Book Distributors, Dehra Dun, pp. 104-138.
- [2]. A.K. Pandey., S. Gayathri, R.C. Rajak and S.K. Hasija, (1996) Possibilities, problems and prospects of microbial management of *Parthenium hysterophorus* in India. In: Perspectives in Biological Science (V.Rai, M.L.Naik & C. Manoharachary eds.), School of life Sciences, Pt. R.S. Shukla University, Raipur , pp. 253-267
- [3]. A.K. Pandey., R.C. Rajak., S. Gayatryi and S.K. Hasija, (1997) Preliminary Assessment of *Colletotrichum dematium* as a mycoherbicide against *P. hysterophorus* L In: Achievements and Prospects in Mycology and Plant Pathology.(S.S. Chahai, L.B. Prashar, H.S. Randhwa & S. Arya eds). International Book Distributors, Dehradunpp.59-85.
- [4]. A.K. Pandey., Jaya Singh and M.A. Varghese (1999). Effect of Rhizobacteria on *Parthenium* PCIN **31**:1-6
- [5]. A.K. Pandey., Jaya Singh, G. M. Shrivastava and R. C. Rajak. (2003) Fungi as herbicides: Current status and future prospects In: Plant Protection: A Biological Approach (Trivedi P.C. ed) Aavishkar Publishers, Distributors Jaipur, India, pp. 305-339.
- [6]. A.K. Pandey., G.M. Shrivastava, A. K. Singh & S. K. Verma (2003) Herbicidal potential of secondary metabolites of *Sclerotium rolfsii* against *Parthenium*: A preliminary observation. *J. Basic Appl. Mycol* **2(1)**: 27-30
- [7]. A.K. Pandey (2001). Microorganisms associated with weeds; opportunities and challenges for their exploitation as herbicides, *Int. J. Mendel*. **17**: 59-62
- [8]. A.K.Pandey, Poonam Chandla and R. C. Rajak, (2002) Herbicidal Potential of Secondary metabolites of some fungi against *Lantana camara* L. *J. Mycol. Pl. Pathol* **32**: 100-102.
- [9]. Archana Pandey and A.K. Pandey (2005) Preliminary assessment of *Colletotrichum dematium* as a potential mycoherbicide against *Hyptis suaveolens*. *J. Basic Appl. Mycol.*, **4**, 21-22
- [10]. Archana Pandey., A.K. Pandey and R.C. Rajak (2002) Mycoherbicidal potential of *S.rolfsii* against *Hyptis suaveolens* L Poit J. *Trop. Forestry*. **18 (1&2)**: 96-98
- [11]. D.F. Bateman and S.V. Beer (1965). Simultaneous productions and synergistic action of oxalic and polygalacturonase during pathogenesis by *Sclerotium rolfsii*. *Phytopathol* **55**: 204 – 211
- [12]. G.J. Kovics, A.K. Pandey and M.K. Rai,( 2005) *Phoma Saccardo* and related genera: Some New Perspectives in Taxonomy and Biotechnology In: Biodiversity of fungi: their role in human life (Eds. S.K. Deshmukh & M. K. Rai), Science Publishers Inc. Enfield (NH), USA. Plymouth, UK, pp. 129-153.
- [13]. H.L. Walker and G.E.Templeton (1978) In vitro production of a phototoxic metabolite by *Colletotrichum gleiosporoides* F.sp *Aeschynomene* *Plant Sci. Letters*. **13**: 91-96
- [14]. H.K. Abbas., C.D. Boyette H.K. Abbas., C.D. Boyette and R.C. Hoagland., *Phytoprotect*. **76**:17-25 (1995).
- [15]. Boyette, and R.C. Hoagland (1995). Phytotoxicity of *Fusarium*, other fungal isolates and of the phytotoxins fumonisins, fusaric acid and moniliformin to jimson weed. *Phytoprotection* **76**: 17-25
- [16]. H.K. Abbas., W.C.A. Glederbloom, M.K. Cawood and W.T. Shier (1993). Biological activities of fumonisin mycotoxins from *Fusarium moniliformin* ,in jimson weed.(*Datura stramonium*L.) and mammalian cell cultures. *Toxicon*. **31**: 345-353.
- [17]. H.K. Abbas., C.D. Boyette and R.C. Hoagland., (1995). *Phytoprotect*. **76**:17-25
- [18]. Knox, Jai, Anita Dass, Manjula Thomas and M.S. Paul (2006). Management of *Parthenium hysterophorus* L. through Atrazine with *Cassia uniflora* extract. *Ann. Pl. Protec. Sci.* **14** (2): **31**:1-6
- [19]. P. Sharma. S. R. Sharma and Sindhu (2004) *Indian Phytopath.* **53 (3)**: 315-318
- [20]. Riti Thapar., A.K. Singh, Archana Pandey and A.K. Pandey (2002) Bioactivity of CFCF of *Curvularia lunata* in *Parthenium hysterophorus* L., *J.Basic Appl.Mycol.* **1**:126-129.
- [21]. O.H.Lowry., Rosebrough, A.L. Fan and R.J. Randa (1951) Protein measurement with the follinphenol reagent. *J. Biol. Chem.* **193**:265-275

- [23]. R.K. Grover and B.L. Chona, (1960). Comparative studies on *Sclerotium rolfsii* sacc. And *ozonium texanum* neal and werster var. Parasiticum Thirumalachar Indian Phytopath., 13; 118-129
- [24]. Rekha Shukla and A.K. Pandey (2006) Maximization of production of oxalic acid from *Sclerotium rolfsii*, a mycoherbicidal agent against parthenium. Ann. Pl. Protec. Sci, **14(1)**: 202-205
- [25]. R.S.Winder and K. Watson (1994) Phytoprotection. **74(1)**: 19-23.
- [26]. S. Joseph., S. Lal and A.K. Pandey (2002) Preliminary evaluation of herbicidal potential of *Streptomyces* WC#150 against *Lantana camara*. Ann .Pl. Protec. Sci. **10**:134-136
- [27]. Sanjai Saxena and A. K. Pandey (2001) Microbial metabolites as ecofriendly agrochemicals for the next millennium. App. Microbial Biotechnol. **55**: 395-403
- [28]. Sanjai Saxena, A.K. Pandey and R.C. Rajak (2001) Potential of an idiolite from *Alternaria alternata* as biorational agrochemicals In: Frontiers in Fungal Biotechnology and Plant Pathogen Relationships (C. Manoharachary, G. Bagyanarayana, B. Bhadracharya, K.Satyaprasad, B.N. Reddy & A. Nagmani eds.). Allied Publishers, Hyderabad, pp.22-28
- [29]. Sanjai Saxena and A. K.Pandey (2002) Evaluation of an indigenous isolate of *Alternaria alternata* (LC#508) for use as a mycoherbicide for *Lantana camara* L. Crop Protection., **21**: 71-73.
- [30]. Sanjai Saxena, A.K. Pandey and R.C. Rajak (2000). Preliminary evaluation of fungal metabolites as natural herbicides for the management of *Lantana camara*. , Indian Phytopath. **53**: 490-493